

'Controlled Hypotension'
in Anesthesia and Surgery

Publication Number 283
AMERICAN LECTURE SERIES®

A Monograph in
The BANNERSTONE DIVISION of
AMERICAN LECTURES IN ANESTHESIOLOGY


Edited by
JOHN ADRIANI M D
Director Department of Anesthesia
Charity Hospital
New Orleans Louisiana

"Controlled Hypotension" in Anesthesia and Surgery

By

DAVID M. LITTLE, JR., M.D.

*Department of Anesthesiology
Hartford Hospital
Formerly Attending Anesthesiologist
Grace New Haven Community Hospital
Assistant Clinical Professor of Anesthesiology
Yale University School of Medicine*



CHARLES C. THOMAS • PUBLISHER
Springfield • Illinois • U.S.A.

Publication Number 233
AMERICAN LECTURE SERIES

A Monograph in
The BANNERSTONE DIVISION of
AMERICAN LECTURES IN ANESTHESIOLOGY

Edited by
JOHN ADRIANI M D
Director Department of Anesthesia
Charity Hospital
New Orleans Louisiana

To

M S L.

Whose normotension has made
this small monograph possible

CHARLES C THOMAS • PUBLISHER

BANNERSTONE HOUSE

301 327 East Lawrence Avenue, Springfield, Illinois U S A

Published simultaneously in the British Commonwealth of Nations by

BLACKWELL SCIENTIFIC PUBLICATIONS LTD OXFORD ENGLAND

Published simultaneously in Canada by

THE DYERSON PRESS TORONTO

This monograph is protected by copyright No
part of it may be reproduced in any manner
without written permission from the publisher

Copyright 1956 by CHARLES C THOMAS PUBLISHER

Library of Congress Catalog Card Number 55 11242

Printed in the United States of America

Contents

<i>Chapter</i>	<i>Page</i>
I INTRODUCTION	3
II BLEEDING DURING SURGERY	5
Non anesthetic Factors	5
Anesthetic Factors	7
III THE CONTROL OF BLEEDING DURING SURGERY	11
IV INDUCTION OF HYPOTENSION	16
V POSTURAL ISCHEMIA	25
VI REDUCTION OF CIRCULATING BLOOD VOLUME	30
Arteriotomy	31
Application of Negative Pressure to the Lower Extremities	37
VII REDUCTION OF PERIPHERAL RESISTANCE CONDUCTION ANESTHESIA	38
Total Spinal Block	41
Epidural Block	46
VIII REDUCTION OF PERIPHERAL RESISTANCE GANGLIONIC BLOCKADE	48
The Methonium Compounds	48
Arfonad	55
Pendiomide	59
IX PHYSIOLOGICAL CONSIDERATIONS	61
The Level of the Blood Pressure	61
Effects on the Cardiovascular System	63
Effects on the Brain	67
Effects on the Liver	70
Effects on the Kidneys	71
The Use of Vasopressor Agents	73
X FATAL AND NON FATAL COMPLICATIONS	75
Complications	75
Mortalities	80

"Controlled Hypotension"
in Anesthesia and Surgery

<i>Chapter</i>	<i>Page</i>
XI CONTRAINDICATIONS	88
XII INDICATIONS	94
Surgical Indications	96
Anesthetic Indications	98
Medical Indications	99
XIII REQUIREMENTS	101
XIV BIBLIOGRAPHY	106
INDEX	137

I

Introduction

THE CONTROL of gross hemorrhage during surgical intervention, and the rapid, adequate replacement of all blood that is lost, constitute fundamental tenets of sound surgical practice. Such control is not always accomplished with facility. Hemorrhage is occasionally so profuse as to endanger the life of the patient despite the most heroic remedial measures. An even more complex problem is the persistent vascular ooze which often renders operation both tedious and difficult and which may amount to a significant loss of circulating blood volume in the course of a comparatively short period of time. Furthermore during certain operative procedures this bleeding need not even be great in quantity to jeopardize surgical success. For example, a single drop of blood during the critical stages of the Lempert fenestration operation may obscure the entire operative field and negate a satisfactory result.

Within the past few years these problems of excessive bleeding during operation have been the subjects of numerous studies both in this country and abroad, and one solution that has been offered consists of the intentional production of hypotension to achieve a relatively bloodless operative site. This concept, while by no means new has recently received widespread notoriety, and is itself sufficiently revolutionary in its character, intriguing in its possibilities, and dangerous in its implications as to warrant detailed consideration.

It would be clearly premature at this time to attempt the proper evaluation of these techniques or to estimate

Prudence does not go behind nature and ask whence it is. It takes the laws of the world whereby man's being is conditioned as they are, and keeps these laws that it may enjoy their proper good. It respects space and time, climate, want, sleep, the law of polarity, growth and death. On the other hand, nature punishes any neglect of prudence.

RALPH WALDO EMERSON

II

Bleeding During Surgery

BLEEDING during surgery occurs only when blood vessels, of whatever type or size, are cut. Normal blood vessels in the absence of pathological processes involving their walls do not ordinarily permit the gross exit of whole blood from their confines. The paramount cause of bleeding during operation therefore is due to the surgical intervention itself. Granted however that this is so, there are nevertheless certain other factors of importance in the causation of bleeding which becomes *excessive* during the operative period. These have been thoroughly reviewed by Gillies who has classified them into two groups according to whether they are anesthetic or non anesthetic in origin (160).

Non anesthetic Factors In the group of non anesthetic factors (Table I) are those specific pathological conditions which prolong coagulation and bleeding times due to vascular deficiency, prothrombin deficiency, platelet deficiency, thromboplastin deficiency or fibrinogen deficiency. Such dyscrasias if untreated pre-operatively are capable of producing varying degrees of abnormal vascular ooze which may become at least troublesome and at times truly dangerous.

Other pathological processes particularly those involving the heart or the lungs may also be a source of increased bleeding in the incised tissues when accompanied by venous congestion and an increased venous pressure. Diseases that produce a high degree of toxicity with concurrent pyrexia and an increased basal metabolic rate

their ultimate position in anesthetic and surgical practice Too much clinical experience has been amassed over too short a period of time to allow the necessary perspective and too little fundamental investigation into the concurrent, profound physiological alterations has been undertaken to provide the essential knowledge Furthermore the subject is too charged with emotional tension on both sides of the question concerning the absolute propriety of these techniques On the one hand there can be no doubt that induced hypotension for the production of a relatively bloodless operative field when wisely employed in carefully selected cases for a rational indication, may prove markedly beneficial in the individual patient Brunschwig has stated that, hypotensive anesthesia may prove to be one of the most important developments in the refinement of operative technic for complicated surgical procedures that has been made in recent years (68) On the other hand there is accumulating a body of evidence which indicates that drastic reduction of blood pressure in order to decrease bleeding carries a risk greater than that normally borne by the patient during surgery Armstrong Davison a pioneer in the development of the hypotensive technique has revised his original position on the subject and asserts that, the facilitation of the surgical procedure is certainly one of the main duties of the anaesthetist but new measures employed to this end must increase the safety of the patient and not be the means of exposing him to unnecessary hazards (99)

Final evaluation of controlled hypotension —be it acceptance or rejection and probably somewhere between the two—must await a precise knowledge of the hazards involved in its application to the human organism, and a careful deliberation upon those hazards in relation to the advantages that can be obtained by its utilization in the individual patient

large amounts of blood from the tissues and structures in the operative area and thus tend to decrease the degree of vascular ooze and hemorrhage

Anesthetic Factors In the group of anesthetic factors (Table II) that must be considered as influencing the

TABLE II
FACTORS INFLUENCING BLEEDING DURING OPERATION

B Anesthetic Factors

1 Agents

- a Effect on bleeding and clotting times
- b Effect of depth of anesthesia
- c Vasodilatation of direct nerve block

2 Techniques

- a Resistance to respiration
 - b Increased intrapulmonary pressure
 - c Carbon dioxide accumulation
-

amount of bleeding during surgical procedures, Gillies has directed attention toward both the agents themselves and the techniques by which those agents are administered (160). He has pointed out the fact that although almost all general anesthetic agents have been blamed in the past for excessive bleeding which occurred during operation yet none of those agents in themselves have any great effect upon the bleeding or clotting times of the blood. Coagulation time is prolonged very slightly during nitrous oxide anesthesia, reduced somewhat during anesthesia produced by ether, chloroform and ethylene, and unchanged during anesthesia occasioned by cyclopropane or the intravenous barbiturates.

On the other hand and of much more importance is the fact that almost all these anesthetic drugs do produce vasodilatation especially of the capillaries of the skin during light planes of general anesthesia. This vasodilatation has been attributed to an initial inhibition of the hypothalamic centers causing a reduction of constrictor

lead to excessive vascularity and thus to excessive bleeding during surgical intervention

Profuse bleeding is most apt to occur in the course of certain specific operative procedures both because of the nature of the tissues and structures being operated upon as well as the pathological processes involving those tissues

TABLE I
FACTORS INFLUENCING BLEEDING DURING OPERATION

A Non-anesthetic Factors

- 1 Deficiencies in the clotting of blood
 - 2 Cardio-pulmonary disease
 - 3 Toxemia Pyrexia
 - 4 Vascularity of tissues
 - 5 Posture
-
-

and structures Thus both glandular tissues and large bodies of muscles are highly vascular, and operations involving them may entail significant blood loss certain intracranial interventions (ligation of congenital aneurysm or very vascular tumors) thyroidectomy intrathoracic operations mastectomy laminectomy, operations on the hip or shoulder joints, and some gynecological procedures are examples of the types of surgery during which excessive bleeding may be encountered because of the nature of the operative site itself

Perhaps the most important non anesthetic factor influencing the amount of bleeding during operation however is the posture of the patient particularly as regards the site of operation in relationship to the dependent portions of the body If the area of operation is dependent in relation to the rest of the body venous engorgement occurs and there is increased vascular oozing into the surgical field. On the other hand when the patient is so positioned that the field of operation is maintained superior to the rest of the body gravitational venous drainage will remove

As will be noted subsequently, however, the production of an ischemic operative field is scarcely a practicable or justifiable solution to the problem of excessive bleeding during surgery if the price to the patient must be a profound generalized cellular intoxication (90)

Drugs employed as adjuvants to anesthesia for the purpose of facilitating muscular relaxation, such as d-tubo curarine decamethonium, or succinylcholine have only weak effects upon ganglionic synapses, except perhaps in truly massive doses. As a result both vasodilatation and active circulatory reflexes combine to cause brisk bleeding during anesthesia supplemented with muscle relaxants, unless of course the general anesthesia is carried to very deep levels.

Even local anesthetic agents will produce vasodilatation with the single exception that when they are employed for local infiltration and a vasoconstrictor is added to prevent toxic absorption vasoconstriction ensues and a decreased amount of bleeding is encountered. When used for direct nerve block, however vasodilatation will occur due to paralysis of the vasoconstrictor fibers of the nerve trunk.

At least as important as the anesthetic agents employed, and in certain instances of far more importance, are the methods and techniques by which general anesthesia is administered. This is most particularly true of the respiratory factors involved in modern methods of anesthetic administration. Increased resistance to respiration, whether due to obstruction of the airway or to increased friction within a closed anesthetic circuit causes an increased venous pressure venous congestion and increased vascular ooze.

Similarly the use of assisted or controlled respirations entail the use of increased intrapulmonary pressures which reduce the intrapleural negative pressure or even make the latter positive lift the thoracic wall and force

tone (160), but it is not a sufficient degree of vasodilatation to lower the blood pressure. Therefore, since the circulatory reflexes are active in light planes of anesthesia this vasodilatation will lead directly to increased bleeding from the surgical wound.

Under such circumstances hemorrhage may be brisk except during anesthesia with those agents which produce a relative hypotension from the very outset, such as chloroform and the barbiturates. Chloroform reduces the blood pressure from the beginning of its administration by a combination of vagal and dilator effects and it is this immediate hypotension that is a large factor in decreased blood loss associated with anesthesia by that agent so much appreciated by the surgeons of an earlier generation (51). The barbiturates have a marked depressant action on the hypothalamic centers controlling autonomic activity and thus produce a hypotension albeit a transient one during the induction of anesthesia. A bit later in the anesthesia, however, the respiratory depressant effects of the barbiturates become more important and offset the decreased tendency toward bleeding attributable to this initial hypotension (128).

A quite different situation pertains during the conduct of *deep* general anesthesia regardless of which agent may be employed to produce anesthesia (42, 517). The vasodilatation seen during light planes of anesthesia persists but instead of being associated with those factors which serve to increase blood flow and therefore active bleeding (i.e., vasodilatation undiminished blood pressure, normal cardiac output and active circulatory reflexes) the vasodilatation during deep anesthesia is accompanied by a depression of the pressor reflexes and a fall in blood pressure. Excessive bleeding therefore can be prevented by maintaining deep levels of general anesthesia, the amount of bleeding varying inversely with the depth of anesthesia.

III

The Control of Bleeding During Surgery

THE PROBLEM of excessive bleeding during surgical interventions is scarcely a new one and for many years both surgeons and anesthetists have applied a variety of different methods in an attempt to achieve hemostasis. For the most part these approaches have been of a local nature, confined to the operative site itself or to that restricted portion of the body immediately surrounding the operative area (Table III).

The use of tourniquets and Esmarch bandages for operations upon the extremities has been a well known maneuver for a very long time. The application of vasoconstrictor substances to the tissues of the operative site has also been employed to provide hemostasis but the utility of this procedure has been somewhat restricted both by the danger of toxic absorption and by the problem of reactive vasodilatation and therefore reactionary hemorrhage. Cautery, chemical agents of an escharotic nature, and more recently hemostatic sponges have proven extremely useful in controlling vascular ooze in fairly limited areas of the operative site. The local application of both heat and cold have been employed for the achievement of hemostasis in the past, and it seems evident from the recent advances in cardiac surgery that hypothermia will gain prominence for the control of hemorrhage in that type of surgery although as a generalized rather than as a local measure and for a quite different rationale than

the diaphragm downward these maneuvers have the tendency to hinder venous return and cardiac filling rather than to facilitate them as does spontaneous respiration and venous ooze is increased

Finally the failure to excrete adequate amounts of the carbon dioxide produced by the patient is a potent cause of increased bleeding during surgery. Uncompensated apnea decreased respiratory exchange due to subapneic doses of respiratory depressant general anesthetic agents or muscle relaxant drugs or inefficient carbon dioxide absorption for numerous technical reasons in closed system methods of inhalational anesthesia all cause a rise in blood pressure and vasodilatation. The importance of the impedance of carbon dioxide excretion as a cause of increased bleeding during surgery has been emphasized recently by the perfection of non rebreathing techniques and the suggestion that they may play a most important role in decreasing vascular ooze during operation (70)

stasis on this basis, of course, is due to the attendant progressive central depression, which leads to hypotension and therefore an ischemic operative field. The gradual realization of the role played by hypotension in producing a bloodless operative site in chloroform, spinal, and deep general anesthetics, has led to the intentional reduction of

TABLE IV
THE HISTORY OF CONTROLLED HYPOTENSION

1946—Gardner—Arteriotomy
1948—Gillies—Total Spinal Block
1950—Davison—Pentamethonium Ganglionic Blockade
1950—Enderby—Hexamethonium Ganglionic Blockade
1951—Guioi and Damousseau—Pendiomide Ganglionic Blockade
1952—Nicholson Sarnoff and Crehan—Arfonad Ganglionic Blockade

blood pressure and the evolution of so-called controlled hypotension during anesthesia and surgery

While it is true that hypotension has often provided a relatively bloodless operative field in the past it has only been sought deliberately for that purpose during the last few years (Table IV). In 1946 Gardner (149) introduced the technique of arteriotomy whereby blood is withdrawn from the patient pre operatively via the arterial route and then replaced in part or in toto by the same manner at the end of surgery. Such pre operative reduction of the total circulating blood volume by the removal of from 500 to 3000 cc of blood serves to lower the arterial pressure to a level between 50 and 90 mm Hg and at the same time produces peripheral vasoconstriction (196) when combined with proper positioning of the patient the net result is a reduction of bleeding at the site of surgery. Arteriotomy thus approximates a state of hemorrhagic shock and for this reason has been thought to exceed the bounds of physiological rectitude.

A more physiological approach to the problem was con-

that upon which its earlier use was based. Arterial compression and even temporary ligation, have been used to control hemorrhage in certain instances of vascular surgery.

Without question, however, the most important of these

TABLE III
METHODS OF HEMOSTASIS

A Local Methods

- 1 Hemostats ligatures and sutures
- 2 Tourniquets and Esmarch bandages
- 3 Vasoconstrictor drugs
- 4 Cautery escharotic chemical agents and hemostatic sponges
- 5 Local application of heat or cold
- 6 Arterial compression or temporary ligation
- 7 Proper posturing of the patient

B General Methods

- 1 Chloroform anesthesia
 - 2 Spinal analgesia
 - 3 Deep general anesthesia
-

approaches to the problem of hemostasis has been the meticulous positioning of the patient on the operating table so that the operative site would be superior to the rest of the body. This maneuver permits gravitational venous drainage away from the operative site into the dependent portions of the body and thus in large measure serves to prevent venous ooze.

It seems possible, in retrospect, that the surgeon's preference in the past for chloroform and spinal anesthesia was due as much to the ischemic operative fields which resulted from the hypotension accompanying these forms of anesthesia as it was to the profound muscular relaxation generally cited as being the basis for his choice of anesthetic procedure. It has also been recognized for some time—and even advocated on a few occasions—that excessive bleeding during surgical intervention could be prevented by recourse to very deep levels of general anesthesia. Hemo-

proaches It might be defined as being a combination of

- 1 The deliberate induction of hypotension

- 2 The employment of Dyle's postural ischaemia

This is in fact, the definition which has come to be accepted in common medical parlance and which will be employed throughout this monograph

tained in Gillies classic studies, in 1948 on *total spinal block* (175) In this method, the blood pressure is reduced to 70 mm Hg, or below, by the use of spinal anesthesia to achieve paralysis of the entire efferent sympathetic outflow, with lesser degrees of sensory and motor blockade so that the muscles of respiration and the medullary centers remain relatively unaffected Gillies has admitted the physiological deviations of even this technique (161), but he has assumed that the arteriolar dilatation so produced guarantees normal oxygenation of the tissue cells, and that the decreased cardiac output inherent in the technique is compensated for by the decreased peripheral resistance and therefore the decreased work of the heart Proper posturing of the patient is again an essential feature for the control of bleeding as it is indeed in all methods of 'controlled hypotension'

The introduction in 1948 of easily injectable ganglionic blocking agents in the form of the methonium compounds (32-362) led to the widespread usage of hexamethonium and pentamethonium to lower the blood pressure to systolic levels of 50 to 60 mm Hg and thus control excessive vascular ooze in the operative field (98-120) More recently, arfonad (344), a thiophanium compound and to a lesser extent, pendiomide, (188) a diethylenetriamine derivative have made possible more readily controllable forms of ganglionic blockade for the purpose of reducing excessive bleeding during operation again employed in conjunction with proper positioning of the patient

A number of other less widely used drugs and techniques have been applied in similar manner to the problem of excessive bleeding during surgery and so-called controlled hypotension is therefore a generic term which includes the total of such diverse clinical ap

less fixed entity under normal physiological conditions can be varied by artificial means as well as by certain pathological processes and therefore may be utilized for the production of controlled hypotension

An intentional decrease of the cardiac output has been

TABLE V
FACTORS CONTROLLING BLOOD PRESSURE

-
- | | |
|---|----------------------------|
| 1 | Cardiac output |
| 2 | Peripheral resistance |
| 3 | Circulating blood volume |
| 4 | Viscosity of the blood |
| 5 | Elasticity of the arteries |
-

sought only by those few workers employing procaine amide (7 311 315 360), a technique of questionable virtue. A decreased cardiac output will of course produce

TABLE VI
FACTORS CONTROLLING CARDIAC OUTPUT

-
- | | |
|---|-------------------------|
| 1 | Venous return |
| 2 | Force of the heart beat |
| 3 | Rate of the heart |
| 4 | Arterial pressure |
-

hypotension but it is important to realize that the reverse may be true also and to emphasize that *all* techniques of controlled hypotension are usually associated with a quite unintentional reduction of cardiac output. This would appear to be unavoidable inasmuch as the cardiac output is mainly dependent upon the venous return, the force of the heart beat, the rate of the heart and the arterial pressure itself (Table VI). Obviously any reduction of arterial pressure may be expected under most conditions at least to lead to a concomitant decrease of cardiac output (208). Furthermore venous return is also

IV

Induction of Hypotension

THE SEVERAL factors which combine to maintain the normal arterial pressure include the cardiac output the total circulating blood volume the peripheral resistance, the viscosity of the blood and the elasticity of the walls of the large arteries (Table V) These various factors are in turn subject to control and influence by an enormous number of other physiological phenomena of nervous hormonal, and physical origin but for the purposes of this discussion the five basic factors mentioned are of sufficient concern

Of these the latter two—the viscosity of the blood and the elasticity of the large arteries—are fairly constant and not easily altered for the purpose of inducing hypotension It is true that the viscosity of the blood fluctuates markedly during surgery changing with variations in the chemical composition the gas content the temperature the colloids and the suspended corpuscles but these changes are not amenable to deliberate alterations with any degree of control The factors of importance for the reduction of the blood pressure in the techniques of controlled hypotension therefore are the cardiac output, the peripheral resistance and the total circulating blood volume

When the blood volume remains constant arterial pressure is to a great extent the product of the cardiac output and the peripheral resistance and these particularly the latter can be varied by devious maneuvers for the intentional reduction of the arterial pressure The quantity of the total circulating blood volume itself, while a more or




Figure 1 Regulation of Arterial Tone VMC vasomotor centers HC higher centers cerebral cortex hypothalamus RC irradiation from respiratory center CO_2 central action of CO_2 O_2 lack central action of O_2 lack CB carotid body (chemoreceptors) CS carotid sinus (pressoreceptors) VA venae cavae and right auricle (pressoreceptors) PA pulmonary artery (pressoreceptors) TS thoracic-splanchnic arteries (pressoreceptors) AB aortic bodies (chemoreceptors) Aortic arch heart (pressoreceptors) CSF cranial and sacral vasodilator fibres PNR posterior nerve roots vasodilator fibres ——— connector and excitor cells of sympathetic nervous system Supr gl suprarenal gland (Modified from Samson Wright *Applied Physiology* 1943 and reprinted from Heymans C *Introduction to the Regulation of Blood Pressure and Heart Rate* Charles C Thomas Publisher 1950 with the kind permission of the publishers)

reduced in hypotensive states, and becomes a further factor in lowering the cardiac output

Hypotension has been induced intentionally, therefore, mainly by creating a disparity between the quantity of the total circulating blood volume and the capacity of the

TABLE VII
THE PHYSIOLOGICAL BASIS OF
THE TECHNIQUES FOR INDUCING HYPOTENSION

-
- | | |
|---|---|
| 1 | Elasticity of the Arteries—relatively unalterable |
| 2 | Viscosity of the Blood—not amenable to control |
| 3 | Decreased Cardiac Output |
| | A Pronestyl |
| | B Unavoidable with other techniques |
| 4 | Reduction of Circulating Blood Volume |
| | A Arteriotomy |
| | B Vacuum applied to lower extremities |
| 5 | Decreased Peripheral Resistance |
| | A Conduction anesthesia i.e. Total Spinal Block |
| | B Ganglionic blockade i.e. Methonium compounds |
-

Y (Modified from Little D M Jr Induced hypotension in anesthesia and surgery *Anesthesiology* 16 320 1955 with the kind permission of the publisher)

vascular bed (Table VII) A reduction of the total circulating blood volume is embodied in the technique of arteriotomy or arterial bleeding followed by retransfusion (6) The *effective* total circulating blood volume is reduced by lowering the peripheral resistance by ganglionic blockade, and then applying a vacuum apparatus to the lower extremities almost an eighth of the total blood volume is trapped in the legs and thighs when suction is exerted on the encased limbs (420)

It has been by a reduction in the peripheral resistance however that the arterial pressure has been most easily and therefore most popularly, decreased for the production of controlled hypotension The peripheral resistance resides mainly in the variably contractile arterioles and to a lesser extent, in the capillaries and veins (Figure 1) The

THE DECREASE OF PERIPHERAL RESISTANCE BY THE INTERRUPTION OF SYMPATHETIC VASOCONSTRICTOR IMPULSES




Figure 2 Interruption of sympathetic vasoconstrictor impulses x

degree of contraction or tone of the arterioles and probably also of the small veins, is maintained and controlled by several factors, of prime importance among which is the continuous stream of vasoconstrictor impulses that arise within the brain and pass down the spinal cord and out over the sympathetic fibers of the peripheral vessels. The peripheral resistance therefore may be reduced by an interruption of this pathway at any point in its course by suppressing the formation of vasoconstrictor impulses, by blocking their pathway to the peripheral vessels, or by preventing the response of the effector cells at the termination of the pathway.

Depression of the autonomic centers which are located in the midbrain will result in a decrease in the number and frequency of constrictor impulses arising in those centers and this effect has been employed extensively in medicine for the control of hypertension by the administration of hydralazine (Figure 2). This is a drug however which has not as yet been found practicable for use during surgical anesthesia.

Depression of the vasomotor center in the medulla oblongata also will suppress the stream of vasoconstrictor impulses. Central vasomotor depression of this type occurs to some degree with most narcotic and hypnotic agents and it is known to occur quite consistently during deep levels of general anesthesia but the deliberate induction of hypotension in this manner has not been considered justifiable because of the generalized cellular intoxication that accompanies such profound central depression.

Reflex inhibition of the vasomotor centers (or stimulation of the vasodilator center) will tend to detract from the effect that vasoconstrictor impulses have upon the peripheral vessels and so will cause an increase in the cross sectional area of the vascular bed and a fall of blood pressure. This is apparently the mode of action of the

cifically to control the hypertensive crises occurring during the removal of adrenal pheochromocytoma

Finally, the response at the termination of the pathway can be prevented by a direct action of such drugs on the

TABLE VIII
TECHNIQUES EMPLOYED IN 27 930 CASES TO PRODUCE
CONTROLLED HYPOTENSION*

Technique	Great Britain and Ireland		U S A		Grand Total of Cases
	Anesthetists	Cases	Anesthetists	Cases	
Arteriotomy	5	206	12	208	414
Total Spinal Block	51	6 922	46	2 655	9 577
Methonium Compounds	166	11 979	115	3 375	15 354
Arfonad*			16	359	359
Pendiomide*			6	63	63
Miscellaneous	35	2 018	14	145	2 163
Totals	255	21 125	209	6 805	27 930

* Questionnaires were sent to the 602 members of the Association of Anaesthetists of Great Britain & Ireland before the general introduction of these compounds into clinical practice

(Reprinted from Little D M Jr Induced hypotension in anesthesia and surgery *Anesthesiology* 16 320 1955 with the kind permission of the publisher)

nitrites the nitrates and certain dehydrogenated ergot preparations, upon the effector cells of the vessel walls themselves. Again, however, none of these has proven to be sufficiently controllable to be employed extensively during anesthesia and surgery

This brief discussion of the physiological basis of the techniques for inducing hypotension would be incomplete without some indication of the relative frequency of their clinical application (Table VIII). Questionnaires were sent to the members of the Association of Anaesthetists of Great Britain and Ireland (197) and to the Diplomates of the American Board of Anesthesiology (198), to ascertain

veratrum alkaloids, although these drugs to date have been used little for controlled hypotension

Sympathetic blockade produced by blocking the preganglionic sympathetic fibers as they emerge from the spinal cord, by one of the techniques of conduction anesthesia has proven to be a most useful method of inducing hypotension during anesthesia and surgery Total spinal analgesia, which has long been known to produce sympathetic paralysis in addition to its effects upon motor and sensory nerves has been employed widely for this purpose Similar results achieved by the same metameric distribution of sympathetic paralysis, have been obtained by the use of epidural analgesia

Sympathetic blockade also may be effected by blocking the sympathetic ganglia themselves Paravertebral sympathetic ganglia blockade (which actually blocks the postganglionic as well as the preganglionic fibers, in some instances) is far too tedious and time-consuming a technique for the purposes of controlled hypotension Sympathetic ganglia blockade however, can be performed with almost indecent ease with the use of such systemically acting drugs as tetraethylammonium bromide or diiodide pentamethonium or hexamethonium halides arfonad pendiomide and others These drugs act to block the transmission across synapses within the autonomic ganglia thereby interrupting the stream of vasoconstrictor impulses and inducing hypotension

Constrictor tone also may be interrupted by blocking the sympathetic nerve endings as they terminate in the walls of the blood vessels by the administration of dibenamine prisol regitine or chlorpromazine These drugs are of uncertain and unpredictable action and have found little use in the techniques of controlled hypotension although they have been employed on rare occasions spe

V

Postural Ischemia

PROPER positioning of the patient upon the operating table can often provide remarkable degrees of hemostasis, and had the techniques of "controlled hypotension" contributed nothing else they would have served at least to reemphasize the value of this salutary maneuver to the present generation of surgeons and anesthetists. In controlled hypotension, lowered blood pressure is combined with elevation of the operative site to cause the elevated part to become pale and ischemic—a situation which Sir Henry Dale has described most aptly as "postural ischemia" (127). It is also true, however, that simple elevation of the area of operation above the rest of the body will in itself effectively lessen the amount of bleeding in the wound.

The promotion of venous drainage is of course the most important factor in this regard. If the surgical wound is dependent in relation to all or much of the body, the venous blood in order to leave the wound must oppose the hydrostatic force of the weight of the column of blood in addition to having to exert an anti-gravitational force. Even when the incision is on a level with the other parts of the body however *venous drainage may be impeded by factors which reduce venous return during anesthesia, particularly venodilatation, diminution of the pump action of the negative pressure within the thorax secondary to decreased respiratory excursion, and because of general muscular paralysis a failure of the intermittent pressure which is normally brought to bear upon the venous blood*

certain facts concerning the clinical practice of 'controlled hypotension'. The results of these questionnaires reported some 27,930 cases of 'controlled hypotension' in anaesthesia and surgery: 15,354 or roughly 55%, were conducted by the intravenous injection of the methonium compounds; 9,577, or 34%, by the induction of total spinal block, which has the advantage of producing surgical anaesthesia over much of the body in addition to inducing hypotension; 414, or 1.5%, by the institution of the somewhat cumbersome technique of arteriotomy; 359 and 63, respectively, by the injection of arfonad and pendiomide which are relatively new drugs; and the remaining 2,163 or approximately 8%, by means of a variety of miscellaneous techniques.

almost entirely upon the nature and site of the surgical procedure, except that due consideration must be given to the dangers of ischemia to vital organs in certain postures. Craniotomies and other neurosurgical procedures




Figure 3 Posture of patient for thoraco-lumbar splanchnicectomy and sympathectomy (Reprinted from Griffiths H W C and Gillies J *Anaesthesia*, 1948 with the kind permission of the authors and the publisher)

in the region of the head and neck are performed in a reverse Trendelenburg position, in a head up tilt of up to 30 degrees or even (posterior fossa explorations) in the sitting position. Many other forms of head and neck surgery (fenestration operation radical neck dissection, mandibular resection) have also required reverse Trendelenburg semi Fowler's or a head up tilt position. Side tilt of the table and some degree of hyperextension have been employed for radical mastectomy while a true lateral jack knife position has been ideal for intra thoracic operations and combined thoraco lumbar splanchnicectomy and sympathectomy (Figure 3). For upper abdominal surgery,

by contraction of the limb muscles (368) When the operative site is superior to the rest of the body, however, these factors combine with the effect of gravity to enhance venous drainage away from the area of surgical intervention and to decrease the amount of bleeding in the wound

It is necessary also to consider the effect of posture upon arterial pressure a factor which generally has been disregarded in the past Enderby has attempted to demonstrate that *local* blood pressure in the wound is reduced 30 mm Hg for every 15 inches of vertical height of the operative site above the level of the heart (125) This is undoubtedly an oversimplification of a very complex hemodynamic situation but it brings out the point that, above the level of the heart gravity opposes the hydrodynamic factor in the arteries, while, conversely, below heart level the hydrostatic and hydrodynamic factors are additive in those vessels

It has been shown that these effects of posture become exaggerated during controlled hypotension conducted by means of a reduction of peripheral resistance because of the marked increase in the capacity of the vascular bed following upon sympathetic blockade Thus, they may be a source of great danger inasmuch as reflex circulatory activity mediated through receptors in the carotid sinuses aortic arch, and the cardio aortic sinus is obtunded under the circumstances of complete sympathetic blockade and positions which impose a marked anti-gravitational factor upon venous return may jeopardize the life of the patient (160) For this same reason however, one can vary the level of the patient's blood pressure by the simple expedient of lowering or elevating a leg an arm or even a hand this method of controlling the degree of the hypotension is a most important one in these techniques

The position in which the patient is placed for operation under controlled hypotension will therefore depend

requirements of cerebral metabolism, and there are a number of workers who refuse to employ controlled hypotension with the head above the level of the horizontal for this very reason. There is the added risk of air embolism in the head up position if large veins are encountered in the field of operation and are inadvertently entered during conditions of venodilatation and hypotension. Other anesthetists have condemned the head-down position because of the possibility that the profound passive congestion of the cerebral vessels during this posture may predispose to thrombus formation (311). The prone and lateral jack knife positions also have been indicted (274), since when the peripheral vascular bed is fully dilated and the inferior vena cava is above the level of its contributory veins venous return may be diminished to a point that is incompatible with continued circulation.

Mention has been made already of the profound reduction of venous return and resultant severe hypotension, that may occur with changes in the patient's posture during complete sympathetic blockade and this point applies post-operatively to the same extent that it does during the operation itself since the closure of the skin incision does not necessarily coincide with the termination of sympathetic blocking activity. Shifts in the patient's posture must be controlled carefully in the post operative period particularly when the legs are lowered from the lithotomy position, or when the patient is moved from the operating table transported on a stretcher or placed into bed.

hyperextension with a raised gallbladder bridge has been used, the hyperextension of the spine in this position probably combining with the use of the bridge to compress the inferior vena cava and so to contribute to the degree of hypotension (124) This is true also of the lateral jack knife position, which has been employed for operations upon the kidney, when the highest point has been just below the thoracic cage The supine jack knife, or a 30 degree Trendelenburg position has provided good working conditions for radical pelvic or abdominal surgery (58) Operations such as repair of the pelvic floor or combined abdominoperineal resection have been difficult problems from the point of view of producing a bloodless field but blood loss can be diminished considerably by the use of the lithotomy position in conjunction with a steep head down tilt (391) A steep head down tilt alone has been used for operations upon the lower extremities but because there appears to be a more sensitive (to ganglionic blockade) sympathetic innervation to the lower extremities than to the upper part of the body better control of bleeding is possible when the site of operation is above the pelvis and legs (440) These examples suffice to show that in general the posture employed for a specific operative intervention conducted under the conditions of 'controlled hypotension' is apt to be closely akin to that which empirical experience had taught many years ago was the most practical position for that particular surgical procedure

There are dangers associated with the positioning of the patient in controlled hypotension of which it is necessary to take cognizance. Certain postures threaten ischemia to vital organs particularly the brain because induced hypotension and gravity may combine to lower the *local* blood pressure (as quite distinct from the blood pressure at heart level) below that necessary to fulfill the

neurosurgical interventions might lessen hemorrhage and so expedite operation, following which the patient could be reinfused with his own blood to reestablish normotension.

Arteriotomy Arteriotomy, as developed by Kohlstaedt and Page (255), appeared to be such a technique, and Gardner therefore induced a relative hypotension in a patient undergoing excision of an olfactory groove meningioma, a large and vascular tumor, to test his theory. A cannula was placed in the dorsalis pedis artery and connected with a reservoir flask, heparin and sodium citrate being employed to prevent clot formation within this system. The pre-operative withdrawal of 1,600 cc of blood served to lower the systolic pressure from 140 to 100 mm Hg, and at the same time increased the pulse rate to 140 beats per minute. It was Gardner's distinct impression that bleeding was greatly reduced during the operative procedure itself and the removal of the tumor facilitated to that extent. At the end of operation prior to the closure of the wound 1,100 cc of the withdrawn blood was reinfused through the arterial cannula; this reinfusion served to raise the systolic pressure from 90 to 120 mm Hg and caused some bleeding from the scalp incision though none from the tumor bed. The patient's convalescence was uneventful. Gardner stressed the fact that the peripheral vasoconstriction which accompanies loss of blood volume was chiefly responsible for the hemostasis and was far more important in this respect than the actual reduction of the arterial pressure.

The use of the technique of arteriotomy and reinfusion to lessen the extensive bleeding that occurs during craniotomy performed for removal of vascular lesions was extended by Hale to include the fenestration operation in which bleeding is not large in amount but rather vexatious and capable of preventing a successful result should

VI

Reduction of Circulating Blood Volume

IN THE COURSE of investigating Seeley's suggestion that the intra arterial infusion of blood or plasma might have advantages over the conventional intravenous route in the treatment of shock (255), Kohlstaedt and Page developed an ingenious method for studying the shock syndrome by means of arterial bleeding and subsequent reinfusion. Essentially, this technique consisted of cannulating the femoral artery in a direction toward the heart and then connecting the cannula to a closed reservoir into which the subject might be bled until the blood pressure had been reduced to the desired level. The blood was mixed with an anticoagulant during its withdrawal and could be reinfused into the femoral artery in order to restore the blood pressure toward normal through the simple expedient of raising the pressure in the closed reservoir by pumping air into it.

Gardner adapted this experimental method of arterial bleeding and reinfusion to the clinical problem of brain tumor surgery (149). He had noted that the hazards and difficulties of excising certain extremely vascular growths abated considerably as the operation wore on: the blood loss increased in amount and the blood pressure fell to hypotensive levels; at that point hemostasis was effected more readily and the remainder of the tumor was removed quite easily. For a number of years he had pondered a technique whereby the withdrawal of blood prior to

CRANIOTOMY GIRL - 14 YRS




Figure 5 Arteriotomy for craniotomy The effect of the withdrawal of 1200 cc of blood upon the pulse and blood pressure is shown (Reprinted from Hale D E *Anesthesiology* 1948 with the kind permission of the author and the publisher)

degree of hemostasis was regarded as satisfactory on an average, this required the withdrawal of 1750 cc of blood over the course of half an hour. If at any time during the bloodletting or the operation itself there was a further fall in arterial pressure blood was returned immediately to the artery from the reservoir under a pressure of 100 to 200 mm Hg until a systolic level of 80 mm Hg or higher had been regained. At the end of the operation, when the need for hemostasis had passed but before the wound had been closed the withdrawn blood was reinfused by both artery and vein (Figure 5). However, since after the removal of a liter or more of blood from the vascular bed, the blood volume tends to increase from its low initial level due to the entrance of tissue fluids into the circulation at a rate which will compensate for the loss of 500 cc of blood in an hour the last 500 cc of withdrawn blood

it occur during the critical maneuver of shaping the fenestra and preparing the skin flap from the auditory canal (196) He used an apparatus which was a modification of that employed by Gardner (Figure 4) and the radial artery—which he ligated at the termination of the procedure—as the site of arterial puncture Blood was withdrawn rather rapidly at the beginning of surgery until a systolic pressure of 80 mm Hg was reached, or until the




Figure 4 Apparatus for arteriotomy The Y shaped glass cannula is secured in the radial artery The lower tubing leads directly to the blood collecting flask The upper tubing leads to a manometer for reading the pressure in the radial artery or in the kelly flask by means of which the blood may be returned to the patient under pressure The small flask and syringe contain a supply of 0.01% heparin solution (Reprinted from Hale D E *Anesthesiology* 1948 with the kind permission of the author and the publisher)

of controllability, both of the arterial pressure as well as the total blood volume (48, 150, 163) that is not possible with most other techniques of controlled hypotension. When reinfusion is necessary, the intra arterial route is by far the most efficient method for raising the blood pressure rapidly (165), particularly in the presence of shock and myocardial ischemia due to decreased aortic pressure (353, 500). Furthermore such reinfusion of the patient's own blood instead of replacing the loss with the blood of one or more donors, not only conserves blood, but also avoids the risks of transfusion reactions or the development of homologous serum jaundice.

Finally, the reduction of both the total blood volume and the arterial pressure has an important advantage for intracranial surgery the tension of the brain is greatly reduced which facilitates retraction during difficult exposures and increases the ease with which congenital aneurysms and other deep seated lesions may be approached (48 295 333).

Despite these advantages however there are several disadvantages to the use of arteriotomy. The major drawback of this technique of course stems from its *modus operandi*—vasoconstriction. The combination of both a reduced total circulating blood volume and the compensatory vasoconstriction which this induces approximates a state of hemorrhagic shock. During such hypovolemic hypotension the ability of the organism to withstand even a minute blood loss is obtunded and the danger of circulatory depression passing into an irreversible stage is constant (48). In this connection furthermore it must be emphasized that blood pressure recordings are not an entirely reliable index of either the state of the peripheral circulation or of the imminence of the phase of irreversibility (517). Even in the absence of an irreversible phase of circulatory depression however there is still the grave

was not administered but was banked and held in reserve for the patient

Other workers have added certain refinements to the technique (27, 48, 229, 295-333) but in essence arteriotomy has been employed in the manner just described. The apparatus has been modified to provide a closed rather than an open system and the heparin has been administered by a slow continuous drip in an attempt to prevent the formation of small thrombi which occurred when heparin was administered by intermittent injections (333). The use of arteriotomy has been confined almost entirely to neurosurgical procedures (i.e., craniotomy) and the fenestration operation (27, 48, 149, 150, 151, 229, 295-333) although recent experimental results indicate that the method may be of great aid in operations upon the open heart and in major hepatic resections (55).


Arteriotomy offers certain well defined advantages as compared to other techniques of 'controlled hypotension'. First and most importantly there is the very obvious hemostatic effect which renders blood loss less severe and thereby facilitates operation to a marked degree. This hemostasis is due apparently to the generalized vasoconstriction neurogenic in origin which has been shown to be the initial response to bloodletting (255) during the brief stage preceding shock. A specific enzyme is believed to be liberated into the plasma which causes this compensatory vasoconstriction (351). Only as the blood volume is reduced further does there result a fall in blood pressure, which then also contributes to the hemostatic effect. For this reason it is unnecessary to lower the blood pressure to dangerous levels in order to obtain a definite hemostasis (150) and bloodless operative fields can be achieved by removing only small amounts of blood (6).

A second great advantage of arteriotomy is the fact that repeated small bleedings and reinfusions permit a degree

of controlled hypotension,' despite its very real advantages in certain respects

Application of Negative Pressure to the Lower Extremities It has been recognized for a number of years that the amount of blood going to the lower extremities can be increased quite markedly by the application of negative pressure to the legs and thighs. It has now been shown that this effect is increased considerably during sympathetic blockade with ganglionic blocking agents (387). Advantage has been taken of this fact to achieve a reduction of the *effective* circulating blood volume by trapping several hundred cc of blood in the legs and thighs during methonium induced hypotension by the use of a negative pressure device which applies pneumatic suction to the lower extremities (Figure 6). This technique has produced controlled hypotension in patients who were resistant to the hypotensive effects of the methonium compounds alone. It has permitted the use of smaller doses of methonium compounds and thereby helped to reduce post operative hypotension, and has allowed a better control over the level of the blood pressure which can be raised or lowered promptly by decreasing or increasing the negative pressure applied to the legs (420-230).

danger of anoxic tissue damage, since the increased peripheral resistance produced by vasoconstriction necessitates higher arterial pressures, in order to maintain cellular respiration, than the low pressures which may obtain during arteriotomy. The fact that this theoretical danger has not materialized during the clinical series that have been




NEGATIVE-PRESSURE DEVICE SHOWING APPLICATION OF CYLINDER TO PATIENT'S LEG

- ✧ Figure 6 Negative pressure device for enhancing methonium induced hypotension (Reprinted from Saunders J W *Lancet* 1952 with the kind permission of the author and the publisher)

reported may be explained in part by the fact that this is the one type of controlled hypotension in which the circulatory reflexes remain active permitting blood to be shunted to vital organs


Certain other disadvantages should be mentioned the equipment is cumbersome and the technique itself is difficult and time-consuming to initiate the dangers of infection and air embolism both exist and have been encountered during the clinical performance of the method and the possibility of clot formation within the apparatus is ever present. These disadvantages together with the phenomenon of vasoconstriction, have served to make arteriotomy the least popular of all the techniques



SYSTEMIC ARTERIAL PRESSURE GRADIENT IN

NORMAL SUBJECT

A: SITE OF MAXIMAL PERIPHERAL RESISTANCE
B: SITE OF TISSUE RESPIRATION



SYSTEMIC ARTERIAL PRESSURE GRADIENT IN THE

TOTALLY SYMPHETOMISED SUBJECT

A: PERIPHERAL RESISTANCE REDUCED
B: NORMAL CAPILLARY PRESSURE SITE OF TISSUE RESPIRATION

VII

Reduction of Peripheral Resistance Conduction Anesthesia

PERIPHERAL resistance resides largely in the variably contractile arterioles and is maintained by the continuous stream of vasoconstrictor impulses arising from the brain stem and carried via the sympathetic nerves to the vessel walls themselves. Interruption of this efferent sympathetic outflow results in arteriolar dilatation and thus effects a reduction of peripheral resistance. The capacity of the vascular bed is increased thereby relative to the total circulating blood volume and this disruption of the finely adjusted balance between the two produces hypotension.

It has been argued that a reduction of peripheral resistance is the most nearly physiological method presently available to induce hypotension for the purpose of achieving a bloodless operative field during surgical interventions (175). The pressure within the vascular tree normally is highest in the aorta and falls progressively as the blood flows through the large and small arteries to the arterioles (267). It is in the vast arteriolar network that the truly pronounced fall in pressure occurs (Figure 7) so that by the

Figure 7 The effect of arteriolar dilatation on the level of blood pressure necessary for tissue respiration (Reprinted from Griffiths H W C and Gillies J *Anaesthesia* 1948 with the kind permissions of the authors and the publisher)

which can be accomplished relatively simply, thus have become the methods of choice for the production of controlled hypotension by means of conduction anesthesia.

Total Spinal Block. The use of total spinal analgesia for surgical interventions has been practised intermittently by bold pioneers since the earliest days following Bier's initial demonstration of spinal analgesia in 1899. In 1900 Morton employed spinal analgesia to accomplish excision of the superior maxilla (334) and operations upon the head under spinal analgesia were reported by Payne as early as 1901 (370). Tait and Cagliari injected at sites as high as the cervical region to obtain total spinal analgesia (469) and Jonnesco utilized this method in over 100 operations above the level of the diaphragm (233). Le Filhatre also practised massive spinal analgesia, employing barbotage to achieve high levels of block (281). The hypotensive effects achieved by the techniques of these early workers were appreciated rather than understood. Pitkin has stated: "In those days (1912) we knew little or nothing about pressure changes and our only common concern was whether we had anesthesia; we used to marvel at the fact that we had such dry wounds" (380).

Koster in the early 1920's was an ardent advocate of total spinal analgesia and employed it intrepidly for such operations as mastoidectomy and thyroidectomy (258, 259). He was thoroughly aware of the marked falls in blood pressure which occurred and approved of them because of the dry wounds that they produced in the non-dependent portions of the body. Bourne and his colleagues have established total spinal analgesia as a valuable method of anesthesia for chest surgery, not least because of the great reduction in the amount of bleeding (56, 190, 191).

It is a matter of present day wonder, in view of the drastic nature of the techniques employed, not that so

time that the blood has reached the capillaries the pressure is only 32 mm Hg. This level of pressure is sufficient to maintain tissue oxygenation and cellular respiration. The hypothesis has been put forward that, if the peripheral resistance is abolished by arteriolar dilatation, very low arterial pressures—32 mm Hg or above to be exact—should be adequate to maintain the tissue oxygenation and cellular respiration inasmuch as the energy of the blood column need no longer be expended in overcoming the frictional surface resistance of the arteriolar bed. In other words the assumption has been made that under conditions of complete arteriolar dilatation and in the presence of a normal blood volume, normovolemic hypotension should be well tolerated by the organism without danger of tissue anoxia (175).

The vasoconstrictor fibers which maintain peripheral resistance belong to the thoracolumbar (sympathetic) division of the autonomic nervous system, and arise from preganglionic cell bodies located in the intermediolateral cell column of all twelve thoracic and the upper three or four lumbar segments of the spinal cord. The axons of these cells traverse the anterior roots, pass to the trunk ganglia as white rami communicantes and are distributed thence to the peripheral vessels either in the outer coats of the large arteries or via the somatic nerve trunks. Blockade of these fibers can therefore be effected by local anesthetic agents acting upon the ventral roots, upon the rami communicantes or upon the sympathetic chain itself.

It has been demonstrated that in order to induce a substantial fall in blood pressure it is necessary to block almost the entire efferent sympathetic outflow (89, 205). Paravertebral and sympathetic chain blocks while theoretically feasible would require too many separate injections and would be too tediously time consuming to be practical. Spinal or epidural (extradural) analgesia both of

paralyzed, and it has been shown that motor block requires the highest and vasomotor block the lowest concentrations of local anesthetic agent, sensory block occupying a somewhat middle position in this regard (Table IX)

Griffiths and Gillies took advantage of these facts to

TABLE IX
CONCENTRATIONS OF PROCAINE REQUIRED TO BLOCK
VARIOUS TYPES OF NERVOUS TISSUE

<i>Type of Nervous Tissue</i>	<i>% of Procaine</i>
Vasomotor Fiber	0.2
Sensory Fiber	0.33-0.5
Motor Fiber	0.5-0.75
Medullary Centers	1.25-1.8

contrive a differential spinal block in which there was motor and sensory paralysis to a level appropriate for the proposed operation but which in addition involved the whole sympathetic outflow and so produced a considerable fall in blood pressure (175). Since a concentration of 0.2% procaine suffices to paralyze preganglionic sympathetic fibers the phrenic nerves were not affected when the anesthetic solution becoming more and more dilute was allowed to diffuse into the cervical region of the cord. They have termed this *total spinal block*, which they differentiated from *total spinal analgesia* in that although the blood pressure was reduced to a level of 70 mm Hg or below by blockade of the entire sympathetic outflow, lesser degrees of sensory and motor block obtained and therefore the muscles of respiration remained relatively unaffected (160, 161, 175).

The basic features of *total spinal block* as evolved by these workers (Figure 8) have consisted of a minimal injection of pentothal to induce hypnosis, the establishment of high spinal block by 150 to 300 mgm of procaine or 1.5

few patients succumbed during these early trials but that any survived at all (175) The surgeon of the era which marked the beginning of the century worked without benefit of the knowledge of the dangers of anoxia or the facilities with which to combat respiratory failure The relatively low mortality that pertained has been ascribed to the fact that anhydrous cocaine was employed and paralyzed the sensory roots but left the motor roots relatively unaffected by the weaker concentrations which reached the upper spinal segments For it has now been well established that death under spinal analgesia results primarily from peripheral respiratory paralysis (131 482) As the level of anesthesia rises into the thoracic region, the intercostal nerves are paralyzed and respiration becomes entirely diaphragmatic and hence slow and of small amplitude Further ascension of the level of anesthesia into the cervical region paralyzes the phrenic nerves and causes respiratory failure

The comportment of the blood pressure during spinal anesthesia has also been the subject of intensive investigation (89 205) Vasodilatation sufficient to effect a reduction of arterial pressure occurs only after the highest thoracic segments have been paralyzed due to the fact that with lesser degrees of blockade intense compensatory vasoconstriction occurs in those parts of the body to which the sympathetic outflow remains intact (343) It has been recognized for some time that it is possible to produce subarachnoid block of sensory fibers that is accompanied by little or no motor paralysis (16) and that when either cocaine (204) or procaine (411) is deposited around a mixed nerve blocking of all fibers does not occur simultaneously The technique of differential spinal block which was developed by Sarnoff and Arrowood (17, 411, 412 413 414) has been employed to investigate the sequence in which the various types of nerve fibers are

vis a tergo (already considerably reduced) be incapable of coping with the additional load. Venous return could be improved, whenever clinical assessment of the state of the circulation dictated, by altering the patient's position (particularly by raising the legs, or tilting the table into Trendelenburg or reverse Trendelenburg, as the situation required) or by abolishing the vasodilatation by the use of vasoconstrictor agents which act peripherally. As the vasoconstrictor paralysis passed off normotension was reestablished. If this did not occur prior to closure a vasoconstrictor (methedrine) was employed to reveal potential bleeding points. The converse problem that of reestablishment of normotension too early in the course of operation may be circumvented by the use of the continuous spinal technique with intermittent injections of the local anesthetic agent being made as required (173-474). Postoperative care must be meticulous with particular attention being paid to the maintenance of a 20 degree head down position for the first eight hours.

As compared to other methods for the production of controlled hypotension *total spinal block* boasts certain well-defined advantages. When the technique is conducted properly and the dosage of the local anesthetic agent for intrathecal administration has been estimated correctly, a profound fall in blood pressure and an extremely ischemic field can be produced in almost all instances. Furthermore this is achieved by paralysis of the sympathetic outflow only whereas blockade produced by chemical ganglionic blocking agents acts upon both parasympathetic and sympathetic ganglia and so introduces other complicating factors. Finally high spinal block is part of the anesthetic technique itself, and so acquires a double usefulness, in that it produces motor and sensory paralysis in addition to the sympathetic blockade. In this connection it must be emphasized that to obtain the fullest benefits from

RECORDINGS OF PULSE RATE, RESPIRATION AND BLOOD PRESSURE OF A PATIENT UNDER TOTAL SPINAL ANALGESIA




Figure 8 Anesthetic record of a patient under *total spinal block* (Reprinted from Griffiths H W C and Gillies J *Anaesthesia*, 1948 with the kind permissions of the authors and the publisher)

to 30 mgm of nupercaine administered intrathecally and maintenance of light general anesthesia with nitrous oxide. The patient was positioned meticulously to permit the operative site to remain superior so that bleeding was minimal and the patient's blood volume remained constant. Since there was generalized arteriolar and post arteriolar dilatation with considerable pooling of blood in the dependent portions of the body emphasis was made of the fact that the anti gravitational resistance to venous return must not be allowed to be too great, lest the arterial

vis a tergo (already considerably reduced) be incapable of coping with the additional load. Venous return could be improved whenever clinical assessment of the state of the circulation dictated, by altering the patient's position (particularly by raising the legs or tilting the table into Trendelenburg or reverse Trendelenburg, as the situation required) or by abolishing the vasodilatation by the use of vasoconstrictor agents which act peripherally. As the vasoconstrictor paralysis passed off, normotension was reestablished. If this did not occur prior to closure, a vasoconstrictor (methedrine) was employed to reveal potential bleeding points. The converse problem that of reestablishment of normotension too early in the course of operation may be circumvented by the use of the continuous spinal technique, with intermittent injections of the local anesthetic agent being made as required (173, 474). Postoperative care must be meticulous, with particular attention being paid to the maintenance of a 20 degree head down position for the first eight hours.

As compared to other methods for the production of controlled hypotension *total spinal block* boasts certain well-defined advantages. When the technique is conducted properly and the dosage of the local anesthetic agent for intrathecal administration has been estimated correctly, a profound fall in blood pressure and an extremely ischemic field can be produced in almost all instances. Furthermore this is achieved by paralysis of the sympathetic outflow only, whereas blockade produced by chemical ganglionic blocking agents acts upon both parasympathetic and sympathetic ganglia, and so introduces other complicating factors. Finally high spinal block is part of the anesthetic technique itself, and so acquires a double usefulness in that it produces motor and sensory paralysis in addition to the sympathetic blockade. In this connection it must be emphasized that to obtain the fullest benefits from

spinal block it is necessary to ensure sympathetic paralysis to a high level even if the sensory and motor blocks are restricted to the segments supplying the operative field since otherwise the compensatory vasoconstriction occurring in the distribution of the unblocked sympathetic fibers may actually increase rather than reduce, bleeding.

Granted, however, that a low head of arterial pressure associated with vasodilatation and a normal blood volume carries less theoretical danger than illusory higher arterial pressures accompanied by vasoconstriction and a reduced blood volume controlled hypotension produced by spinal anesthesia still carries with it those inherent capacities for catastrophe which have become the hallmark of these techniques. Furthermore, the usual disadvantages associated with spinal analgesia such as headache, sepsis, and the like still apply. The technique lacks the controllability that is obviously desirable and found to some degree with certain ganglionic blocking agents (i.e. arfonid see Chapter VIII). Finally when a spinal block has been administered to a sufficiently high level to permit upper abdominal or mid abdominal surgery assisted respirations may be necessitated by a diminution of respiratory exchange, and this maneuver by increasing intrapulmonary pressure and therefore venous back pressure may actually cause an increase in vascular ooze and defeat utterly the objective of the technique.

Epidural Block. Epidural analgesia has been advocated for the control of hypertension in eclampsia and the toxic states of pregnancy on numerous occasions during the past decade but only recently has it been utilized to produce intentional hypotension during surgical procedures (62, 63, 114). Indeed one of the major advantages ascribed to epidural analgesia in the past has been the absence of effect upon blood pressure. This is of course true only if the block is truly segmental and the number of seg-

ments blocked are few in number. However, if the block extends over the lumbar outflow and well up into the thoracic segments, sufficient sympathetic fibers are paralyzed to result in a profound fall in arterial pressure (89, 205).

Bromage has been a leading exponent of this method for producing "controlled hypotension" and has patterned his technique closely after *total spinal block* as developed by Gillies, extradural block being substituted for the subarachnoid block (62, 63). The advantages cited include all those attending spinal block, but without the inherent dangers of headache, meningismus and the like. There remain the disadvantages of the greater difficulty in performing epidural block as compared to the performance of lumbar puncture, the admitted inability to prejudge the effective height of an epidural block with the same degree of accuracy as a subarachnoid block, the grave hazard of massive subarachnoid injection, the possibility of infection of the epidural space and a failure rate of 1.4% (even in obviously expert hands) (63).

One novel method of controlling bleeding in the operative site by the use of epidural analgesia has been the establishment of partial sympathetic blockade, sufficient to induce a mild hypotension, but insufficient to prevent an intense compensatory vasoconstriction in the upper part of the body (343). This combination of generalized hypotension and localized vasoconstriction in the operative site has been exploited particularly in the fenestration operation and other forms of cranial surgery (168).

VIII

Reduction of Peripheral Resistance. Ganglionic Blockade

PERIPHERAL resistance can be reduced by interrupting the continuous flow of vasoconstrictor impulses to the arterioles at any point in their route. Of all the methods for inducing hypotension by the reduction of peripheral resistance the use of ganglionic blocking drugs has proved to be by far the most popular technique in the production of controlled hypotension (see Table VIII), due in large measure no doubt to the fact that the administration of these agents requires merely a simple intravenous injection.

Certain generalizations can be made concerning the use of these drugs. They are not anesthetic agents in themselves although clinically the hypotensive state does seem to be associated with a need for less general anesthesia than normal. These autonomic ganglionic blocking agents paralyze both parasympathetic and sympathetic ganglia thus introducing undesirable complicating factors in certain instances. Finally, the failure rate (that is the failure to produce hypotension) is often high when these drugs are employed for that purpose much higher than that encountered in the use of either arteriotomy or the various methods of conduction anesthesia that are utilized to reduce peripheral resistance.


The Methonium Compounds Pentamethonium and hexamethonium are respectively the five and six carbon

atom homologues of the polymethylene series of methonium compounds (Figure 9), the pharmacological properties of which were described by Piton and Zaimis in 1948 (363). The ten carbon atom homologue, decamethonium,

THE METHONIUM COMPOUNDS

POLYMETHYLENE BIS TRIMETHYL AMMONIUM COMPOUNDS

DECAMETHONIUM



HEXAMETHONIUM



PENTAMETHONIUM




Figure 9 The methonium compounds

which has a potent myoneural junction blocking activity (350) has achieved widespread use for the production of skeletal muscle relaxation during clinical anesthesia (348). Pentamethonium and hexamethonium on the other hand do not block the myoneural junction to any such significant degree but do inhibit transmission of nervous impulses through the autonomic ganglia of the sympathetic and parasympathetic nervous systems (364). These potent and specific ganglionic inhibitors apparently act to paralyze ganglionic transmission by raising the threshold of the ganglion cell to acetylcholine released at the preganglionic nerve endings (365). It is this blockade or inhibition

of ganglionic transmission which is responsible for the major pharmacological properties of penta and hexa methonium vasodilatation fall in blood pressure, and an increase in blood flow and elevated skin temperatures in the extremities (14 15 71, 177, 423, 472), inhibition of intestinal peristalsis, inhibition of gastric secretion and reduction of gastric motility (108, 237, 238), inhibition of salivary secretion (472) and dilatation of the pupils together with some disturbance of accommodation of the eye

Clinical application of these pharmacological properties has been made by the employment of the methonium compounds for the treatment of gastric and duodenal ulcer (237 238 433), peripheral vascular disease (71, 136 137, 146 427) acute ventricular failure (499), intractable hiccup (212, 301 302) toxemias of late pregnancy (330 374 473) and most importantly hypertension (74 93 143 305 307 386, 431 453, 455) The administration of these drugs in the medical therapy of the various hypertensive states has brought to light a number of points of some importance in connection with their use in the production of controlled hypotension in anesthesia and surgery

It is now generally accepted that although both penta and hexa methonium will produce vasodilatation and a fall in blood pressure the six carbon homologue is probably the more predictable and dependable of the two in this regard (74 140 456) however it should be pointed out that not all observers concur in this view (33) It must be recognized that tolerance may develop to these drugs and for this reason it may become impossible to maintain a satisfactory hypotension (142 252, 473) Indeed, individual resistance may obtain from the very commencement of therapy so that a significant fall in blood pressure is never achieved (30)

On the other hand prolonged hypotensive effects may

follow the use of these agents (495), particularly in the presence of kidney disease and impaired renal function, since these compounds are excreted almost quantitatively in the urine by both the experimental animal (513) and man (324). The mechanism of the renal excretion of the methonium compounds was studied by comparing the clearances of these substances with the simultaneous clearance of inulin, and from the methonium/inulin clearance ratios so obtained it was clearly evident that the renal excretion of methonium compounds is due mainly to glomerular filtration with only minimal tubular excretion (512). Therefore, even in the absence of renal disease, methonium induced hypotension may decrease glomerular filtration sufficiently to delay excretion of the drug and thus produce a vicious cycle of prolonged activity (306-308). There is also the further danger of bromism or iodism (217, 287, 361-392, 460) if renal function is abnormal or if large doses of either of these salts are employed over a prolonged period of time; however, this hazard can be circumvented by the use of the bitartrate or chloride preparations (75).

Finally there is the every grave problem of dosage, and especially of initial dosage (30) for the individual variation in response to a given dose of these substances is immense (142, 144-289), varying not only between patients but also in the same patient at different times (499). On the one hand this may make a satisfactory therapeutic response to the administration of these drugs difficult or even impossible to attain (252-289-341) and on the other hand a sudden and extreme hypotension may be induced and result in pathological damage to vital organs (213-454), particularly when vascular sclerosis already exists.

Penta and hexa methonium were employed first in anesthesia as antidotes to the muscle relaxant decamethonium (350) before the hypotensive effects of these com-

pounds were generally appreciated. It became apparent immediately that these drugs produced a profound fall in blood pressure when administered to the anesthetized patient (227) and that they should not be employed as antagonists to myoneural blocking agents for this reason. However, this hypotensive effect was seized upon immediately as the basis for a technique of 'controlled hypotension' (98, 120), and the success achieved thereby led rapidly to the extensive use of the methonium compounds for the medical therapy of hypertension (74 93 143 305 307 386 431, 453, 455).

The administration of methonium compounds to induce hypotension for the reduction of bleeding during operative interventions has now become exceedingly wide spread (18 19, 59 78 79 96 110, 123, 124, 126, 127, 211 214 224 231, 239 248 274 279 285 291 303 315, 340 347 349 373 390 399 435 440 465, 479 491 494 502). A number of different techniques have been developed by the many workers who have employed the methonium drugs to induce hypotension but these techniques differ mainly in details and minutiae. The broad general plan of the majority of these techniques is similar to that illustrated in Figure 10. The anesthesia is induced by the administration of intravenous pentothal intubation is performed (a muscle relaxant usually being employed to facilitate this maneuver) and anesthesia is then maintained with an inhalation agent (nitrous oxide cyclopropane ether or trichlorethylene (98 120 127). The methonium salt is then administered intravenously to produce hypotension. hexamethonium is more consistent in achieving this effect and for that reason enjoys a greater vogue than the pentamethonium homologue (19, 59, 79 127 214 231 239 279 303 491 494 502). Two schools of thought have developed regarding dosage one group of workers feels that a large (50 to 100 mgm) initial dose is

as the clinical conditions warrant (59, 462) The rationale behind this latter technique has been carried to its logical conclusion by a few who have employed hexamethonium in a dilute intravenous infusion (274, 394) A certain group of patients (about 15%) are resistant to the effects of the methonium compounds it is usually young and robust individuals who fall into this category (126) Another, smaller number of patients develop resistance to repeated doses and it is for this reason that some workers feel that the initial dose must be large enough to achieve the desired fall in pressure, for the required duration of time without recourse to repeated administration (127)


Following the initial fall in blood pressure the patient is positioned so that the operative site will be superior to the other portions of the body, and this maneuver usually provides a further reduction of the systolic pressure due to pooling of the blood in the dependent portions of the dilated vascular system (71) Operation is then begun Bleeding is usually minimal, but all the blood that is lost must of course, be replaced *in toto* Operation generally proceeds with dispatch and the operative time is diminished to a considerable extent There is a marked difference of opinion concerning the desirability of employing vasoconstrictor agents at the end of operation for the purpose of raising the pressure and revealing potential bleeding points prior to closure of the wound (197, 198) the argument against the use of these drugs is that clots and thrombi which have formed in the cut dilated vessels will be retained *in situ* if the vessels gradually regain tone and contract down about the formed clots whereas sudden increases of intravascular pressure produced by vasopressor drugs will tend to blow out such clots (see Chapter IX)

Post operative care must be meticulous perhaps more so with this technique than with other forms of controlled hypotension for the methonium compounds may have

prolonged effects upon vessels, making changes in the patient's position dangerous for several hours after their administration, even though the blood pressure has returned to normal when the patient is in the horizontal position (71)

Arfonad Arfonad, a thiophanium compound with po

ARFONAD



4,3,4 (1',3') **DIBENZYL 2'-KETO-IMIDAZOLIDO** 1,2-TRIMETHYLENE
THIOPHANUM d CAMPHOR SULFONATE

Figure 11 Arfonad

tent ganglionic blocking activity (384) has afforded the anesthesiologist a more controllable method of inducing hypotension by means of reduction of the peripheral resistance than can be attained by either conduction anesthesia or the use of the methonium compounds (344). Chemically this drug is d 3 4 (1 3 dibenzyl 2 ketoimidazolido) 1 2 trimethylene thiophanium d camphor sulfonate (Figure 11). It apparently has direct vasodilator

properties as well as potent ganglionic blocking activity (300) and single intravenous doses of 0.1 or 0.2 mgm per kgm of body weight will produce a prompt but evanescent depressor response of the arterial pressure (417). Thus it is possible to employ the drug by a continuous intravenous infusion technique to provide graded and rapidly reversible falls in arterial pressure. It has been administered in this manner in the treatment of acute pulmonary edema in hypertensive crises (95, 410, 415, 416, 417, 418, 419) to control sympathetic overactivity in ulcerative colitis (511), and in the management of hypertensive complications of pregnancy (22). The mechanism responsible for arfonad's very brief action is poorly understood, but there is evidence that at least 30% of the compound is excreted in the urine, apparently unchanged, with great rapidity (156).

The use of arfonad in controlled hypotension during surgical interventions has been by the continuous infusion technique, and has provided a degree of control over the level of the blood pressure on a minute to minute basis that has not been obtainable with other methods of induced hypotension. The anesthetic techniques and management have been very similar to those employed when hypotension is achieved by the administration of the methonium compounds (4, 28, 76, 102, 113, 218, 251, 286, 310, 329, 400, 401, 402, 403, 437, 489).

One innovation made possible by this drug, however, has been the development of a modified technique of controlled hypotension (286). This modified concept has been based upon two important facts. First, that the level of the systolic arterial pressure necessary to produce a relatively bloodless operative field varies tremendously between individual patients, and therefore the exact level to which the pressure must be lowered cannot be foretold with any degree of accuracy prior to actual operation. This knowledge can be obtained only after suitable pos-





Figure 12 Anesthetic chart of modified technique of arfonad induced hypotension (Reprinted from Little D M Jr Hampton L J and Grosskreutz D C *Surgery* 1954 with the kind permission of the publisher)

turing of the patient and direct observation of the bleeding—or lack of it—in the operative field itself. Second, this level of optimal pressure is often much higher than the dangerously low pressures (50 to 60 mm Hg) often advocated in the past. Indeed, a significant reduction in

PENDIOMIDE

CIBA 9295



PENTAMETHYL-DIETHYL-3-AZA-PENTANE-1,5-DIAMMONIUM DIBROMIDE

Figure 13 Pendiomide

blood loss is frequently noted at pressures of 90 to 100 mm Hg, and this may represent only a 10 to 20 mm Hg decrease in pressure from the preoperative normal. Under such circumstances, the benefits of controlled hypotension become available to the patient at less risk of major sequelae.

In this modified technique (Figure 12) following the completion of anesthetic induction and proper posturing of the patient, the arfonad infusion in a 0.1% concentration is started and run at a sufficiently rapid rate to evoke a substantial fall in arterial pressure. As the pressure is being lowered, repeated blood pressure readings are correlated with the degree of vascular ooze visible in the wound in order to ascertain the all-important level of pressure

which will provide control of excessive bleeding during operation. The infusion rate is then regulated to maintain this blood pressure throughout that part of the operation during which it is desirable to control the amount of




Figure 14 Anesthetic record of hypotension induced with pendiomide (Through the kind courtesy of Dr C R Stephen)

bleeding. At the end of surgery or in the event of hemorrhage or similar surgical exigency normotension can be restored rapidly by merely discontinuing the infusion.

Arfonad is not quite the panacea for controlled hypotension, however, for even this admirable drug has certain disadvantages such as the development of tachyphylaxis, the occasional onset of tachycardia under its effect and rarely prolonged hypotensive effects.

Pendiomide A third method of chemical ganglionic blockade for the purpose of lowering peripheral resistance

has been by the administration of the diethylenetriamine derivative, pendiomide (188). This drug, chemically $\text{N,N,N',N',3 pentamethyl N N' diethyl 3 aza pentane 1 5 diammonium dibromide}$ (Figure 13) and it has been shown to have marked ability to inhibit the synaptic transmission of impulses through autonomic ganglia (37 38 44 262, 294, 314). Pendiomide has been employed for the reduction of intra ocular pressure in glaucoma (44) for the treatment of eclampsia (44), for the therapy of herpes zoster (346), for relief of bladder tenesmus during cystoscopy (428) for the therapy of acute pulmonary embolism (91 429) and of course, for the production of controlled hypotension during surgery (24 41 80 106 155 187 188 189 235, 254 407 451 478 516).

The anesthetic techniques and management which have been utilized in conjunction with the use of this agent for the production of controlled hypotension have been very similar to those described for methonium or arfonad induced hypotension (Figure 14). A single intravenous injection of pendiomide has an acute effect of five to 10 minutes duration complete recovery occurring in about 30 minutes. Pendiomide is therefore somewhat between the methonium compounds and arfonad as far as duration of activity is concerned and the average dose necessary to achieve an optimal hypotensive effect is between 50 and 100 mgm administered at the rate of 10 mgm per minute. Its advantages include high specificity of action and low toxicity while its major disadvantages are the facts that the dosage is unpredictable and that some patients are resistant to the effects of the drug. It appears to be less controllable than arfonad although more controllable than the methonium compounds.

IX

Physiological Considerations

GILLIES has referred to the techniques of induced hypotension during anesthesia as physiological trespass (161) and there can be no doubt that they frequently do exceed the bounds of physiological rectitude. It has been a source of considerable amazement that so many anesthesiologists and surgeons trained to abhor and combat any deviation from normotension, should so suddenly embrace the reactionary philosophy of deliberate hypotension. A source of even greater amazement has been the fact that they have done so in the face of a glaring lack of fundamental investigation into the profoundly disturbed physiology that is inherent in these techniques. It would seem imperative that those employing controlled hypotension should understand fully the nature of the physiological derangements so produced particularly as to the state of such vital organs as the heart the brain the liver and the kidneys.

The Level of the Blood Pressure The levels of the blood pressure sought (but presumably not always attained) in 27 930 cases conducted under the techniques of controlled hypotension (197-198) are recorded in Table X. It is worth noting that these blood pressures low as they may seem in many instances were in all probability even lower in actual fact. Woodhall and his associates have demonstrated a significant variation between the pressure readings obtained by the arm cuff method and the actual systemic intravascular pressure as recorded by a Sanborn electromanometer from the cannulated radial

artery during controlled hypotension produced by ganglionic blocking drugs. The arm cuff method showed an average fall of 34.6% in blood pressure under these conditions, whereas simultaneous strain gauge recordings indi-

TABLE X
THE LEVEL OF BLOOD PRESSURE SOUGHT
IN 27 930 CASES OF CONTROLLED HYPOTENSION

<i>Blood Pressure Level</i>	<i>Great Britain and Ireland 21,125 Cases</i>	<i>U S A 6,805 Cases</i>	<i>Grand Total 27,930 Cases</i>
60 Mm Hg or Less	4,571	329	4,900
No Fixed Level but Above 60 Mm Hg		968	968
Pre operative Diastolic Level		50	50
60-80 Mm Hg	6,847	2,588	9,435
70-80 Mm Hg	5,727	166	5,893
80-90 Mm Hg	1,392	2,664	4,056
90 Mm Hg	666	3	669
100 Mm Hg	232		232
Not Stated	1,690	37	1,727

(Reprinted from Little D M Jr *Induced hypotension in anesthesia and surgery Anesthesiology* 16:320 1955 with the kind permission of the publisher)

cated an actual fall of 52.9% in radial arterial pressure (507)

Low levels of arterial pressure are undoubtedly seldom physiological when accompanied by vasoconstriction, but a considerable body of evidence has been presented to support the contention that hypotension *per se*, is not necessarily a dangerous phenomenon if brought about by arteriolar dilatation and without loss of circulating blood volume. Phemister and his co-workers have shown that reduction of the systolic blood pressure in rabbits to 25 to 40 mm Hg produced by stimulation of the aortic depressor nerve could be maintained for a far longer time without lethal outcome than an equal reduction of pressure produced by hemorrhage (379). Dale and Laidlaw

found that a carefully adjusted infusion of acetylcholine could produce dilatation of the cat's arterioles and lower the blood pressure to 30 to 40 mm Hg, although the veins remained well filled and the venous return to the heart was normal, on the other hand when a similar reduction of pressure was produced by a slow infusion of histamine, the venous return failed and the blood was found to be in the engorged capillary bed (94) a condition which Moon has described as irreversible shock (328)

It seems evident from these animal investigations, as well as from comparable observations during many thousands of clinical experiences with the techniques of controlled hypotension that the mammalian organism can indeed survive prolonged periods of profound hypotension produced by arteriolar dilatation and undoubtedly can do so under such circumstances far more often than it can survive a similar hypotension produced by either capillary dilatation or hemorrhage. The interesting question arises however as to the effect of such low pressures upon the organism itself when it does survive and the role that these hypotensive levels of blood pressure may play in producing mortalities in those who fail to survive.

Effects on the Cardiovascular System Studies of the pulse rate during operations performed under high spinal analgesia have usually revealed a definite bradycardia (296 297 409) whereas hypotension induced by a ganglionic blocking agent was more likely to be accompanied by a relative (285) or definite (390) tachycardia. The discrepancy between these effects of controlled hypotension upon the heart rate may be explained in part by the fact that the sympathetic blockade of high spinal analgesia evokes the Bainbridge reflex the lowered venous pressure within the great veins producing reflexly an increase in vagal tone a bradycardia and a prolonged diastole (175) whereas hypotension induced by ganglionic blocking

agents is accompanied by parasympathetic blockade in addition to the sympathetic blockade and in some individuals the former may predominate, with a resultant tachycardia

The stroke volume of the heart during operations performed under high spinal analgesia was reduced from an average control level of 57.3 cc to 28.3 cc and the cardiac output and cardiac index fell to 40% of the basal level (296-297-409). Studies of the cardiac output during hypotension induced by ganglionic blockade have shown conflicting results due in large measure to differences in the clinical conditions during the observations, particularly as concerns the degree of hypotension and the posture of the subject during the period of hypotension. In the recumbent position cardiac output in the unanesthetized subject was usually well maintained (145-499). When however such patients were tilted from the horizontal to a foot down position a significant reduction of cardiac output occurred. Cardiac output in surgical patients under conditions of 'controlled hypotension' induced by methonium compounds was reduced 53% on the average and as much as 75% in certain patients (445). Investigation of the cardiac output by ballistocardiography during arfonad induced hypotension likewise showed a diminished stroke volume although the accompanying compensatory tachycardia prevented marked alteration of the minute volume (401).

Measurements of right auricular pressure during high spinal block demonstrated a profound reduction (296-297-409) which would be expected in view of the fact that the venous pressure is greatly reduced and there is a marked diminution in venous return to the heart both after the induction of high spinal analgesia as well as following the administration of ganglionic blocking agents (185-427). Pulmonary arterial pressure was decreased

markedly also, as much as 34% following high spinal analgesia (409) and to an even greater extent after the induction of hypotension with ganglionic blocking agents (162)

The oxygen consumption, in the studies made of high spinal analgesia, decreased an average of 33% but the arteriovenous oxygen difference rose from an average of 4.07 vol % to 6.79 vol %. The oxygen saturation of the arterial blood was not significantly altered in spite of a slight decrease in the arterial oxygen capacity (which was due to an unexplained fall in hematocrit that amounted to 3.3%) (296, 297, 409). The oxygen consumption in the cat made hypotensive by the administration of either arfonad or hexamethonium was likewise perceptibly reduced (369).

The calculated work of the left ventricle was decreased on an average of 85% from the control level (297, 409) and led to a marked reduction of coronary flow in the surgical patient undergoing high spinal block (409). A similar decrease in coronary flow has been found in the experimental animal made hypotensive by either the subdural injection of procaine or the intravenous administration of etamon (115), but more recent studies in anesthetized dogs with renal hypertension (experimental perinephritis) made hypotensive by the intravenous injection of hexamethonium showed that coronary blood flow increased slightly and that coronary vascular resistance decreased during the period of reduced blood pressure (92). A reduction of coronary flow would of course be profoundly significant from the clinical viewpoint and some insight into the matter has been gained from electrocardiographic tracings made during controlled hypotension (72, 195, 467). Intravenous hexamethonium and pentamethonium are capable of producing changes in the EKG characterized by a vacillating pulse rate, extrasystoles, alteration in the site of the pacemaker and other arrhythmias and de

pression of the ST segment and T wave, suggestive of coronary insufficiency (127, 179) Similar changes occur during the conduct of controlled hypotension including depression of the ST segment and the T wave, disappearance of the P wave and shortening of the PR interval (467) but these changes have been dismissed by the proponents of induced hypotension as not indicating myocardial ischemia (230) this is perhaps an unjustifiable assumption

The peripheral blood flow as measured in the digits of the hands and feet is increased under high spinal analgesia despite the hypotension and decreased cardiac output (296 297, 409) Ganglionic blockade with methonium compounds also effected an increase in the mean blood flow of the foot from an average control level of 10 cc /100 cc of foot volume to an average level of 9 ■ cc /100 cc of foot volume (427) This increase in peripheral blood flow following the administration of hexamethonium and pentamethonium to the unanesthetized supine human being (14 71 145) however was not demonstrated in the patient under anesthesia (445)

Splanchnic blood flow following the administration of hexamethonium to the unanesthetized recumbent human subject fell in proportion to the fall in arterial blood pressure but no change was found in splanchnic vascular resistance (388) This is rather surprising in view of the fact that splanchnic vasodilatation has long been thought to play an important role in the production of the hypotensive state This data on splanchnic blood flow has been confirmed by studies during both low and high spinal analgesia in human beings, no significant changes in splanchnic resistance or oxygen consumption being found (338)

The circulation time ■ greatly prolonged following both spinal analgesia and ganglionic blockade produced

by ganglionic blocking agents (162), sometimes to twice the normal rate (145) and it is obvious that this slowing of the circulation may predispose to thrombosis and emboli particularly if the vessels are sclerotic.

Effects on the Brain Since the brain, of all the tissues of the body is the least tolerant of anoxia, there has been considerable speculation over the effects of controlled hypotension upon that organ. Two avenues of investigation have been pursued in this regard the nitrous oxide technique developed by Kety and Schmidt has been employed to measure cerebral blood flow (249) and cerebral function has been measured by both psycho-physical (345) and purely physical (65-425) tests.

Studies of cerebral hemodynamics during induced hypotension in the unanesthetized subject have yielded conflicting results and cannot in any event be interpolated directly into the situation that pertains during controlled hypotension inasmuch as general anesthesia becomes a factor that cannot be ignored in the equation. These investigations have been of great interest however for they bring out several facts of importance.

In the first place it appears that in the supine position cerebral oxygen consumption is maintained near normal whether the induced hypotension is accompanied by a significant decrease in mean cerebral blood flow (331-332-337) or not (92-103) and that the compensatory mechanism responsible for this maintenance of near normal cerebral metabolism includes both a decrease in cerebral vascular resistance (92-103) and a more complete extraction of oxygen from the blood flowing through the brain as measured by an increased arteriovenous oxygen difference.

A second important fact that has been revealed is that when a severe head up tilt of 30 to 40 degrees is superimposed upon hexamethonium induced hypotension (464)

in the unanesthetized subject these compensatory mechanisms no longer suffice to maintain cerebral metabolism and a critical level of cerebral blood flow is reached (between 23.9 and 45.8 cc per minute per 100 Gm Brain tissue with an average value of 35.0 cc per minute per 100 Gm brain tissue) at which the signs and symptoms (i.e. perspiration, cold clammy skin yawning somnolence slurring speech nausea and occasionally, convulsions) of cerebral ischemia become manifest (138). Similarly when hypotension is induced acutely by differential spinal block, there follows a significant diminution of the mean cerebral blood flow (250). Finally and perhaps very obviously, this critical level of systemic blood pressure at which cerebral ischemia occurs is of course much higher in the hypertensive subject (100 to 110 mm Hg mean femoral arterial pressure) (138).

Investigation of cerebral blood flow under the actual conditions of controlled hypotension during operation has been attempted by measuring the cerebral arteriovenous oxygen difference and then calculating the estimated cerebral blood flow (421). These studies were all made with the patient kept horizontal or nearly so and for this reason the results unfortunately shed no light upon cerebral hemodynamics when Trendelenburg or the reverse Trendelenburg positions are employed during controlled hypotension. Hypotension was produced by the use of hexamethonium and the application of pneumatic suction to the patient's legs (230-420). Under these conditions the arteriovenous oxygen difference was found to be 4.8 vol % (less than normal) and the estimated cerebral blood flow was 46.9 cc blood per minute per 100 Gm brain tissue. Since this figure is roughly the equivalent of that found in a healthy adult breathing oxygen it was concluded that cerebral blood flow was certainly adequate to meet the metabolic demands of the brain in the presence

of the decreased cerebral oxygen demand occurring during general anesthesia. It was also concluded that since the arterial blood pressures of these subjects ranged from 40 to 65 mm Hg during the observations that it may be assumed that controlled hypotension at such blood pressure levels does not produce deleterious effects upon the cerebral hemodynamics if the patient remains approximately horizontal.

There is at least suggestive evidence however obtained both from psychophysical and from purely physical tests of cerebral function that cerebral metabolism is not maintained at a normal level during controlled hypotension in which the blood pressure is reduced to low levels and the patient is postured in the head up position.

Nilsson has adapted the flicker fusion frequency test to the post-operative study of 15 patients who had been operated upon under hexamethonium induced hypotension. Six of these 15 patients showed positive tests indicating cerebral injury presumably anoxic in origin and it is most interesting that although the cerebral injuries that were thus demonstrated were not serious five of the six positive tests occurred in patients who had been positioned with the head end of the table raised between 10 to 20 degrees during the operative (and hypotensive) period (345). This indicates that postural ischemia of the brain is a very real danger under these circumstances.

Further evidence in this direction has been supplied by Bromage who employed the electroencephalogram as a measure of cerebral metabolism. It has been recognized for some time that the electrical activity of the brain as recorded by the electroencephalogram becomes abnormal during cellular anoxia: the normal alpha rhythm of 10 cycles per second is replaced by slow delta waves of less than four cycles per second and there is a reduction of the total electrical energy of the cortex. In three conscious

volunteers who received pentamethonium intravenously and then were postured in a foot down tilt, the severe hypotension so produced was accompanied by coma convulsions and profound disturbances of cortical rhythm the normal alpha rhythm being replaced by slow delta waves in irregular combination (65) It is obvious that conclusions drawn from this study must be limited inasmuch as the surgical patient undergoing controlled hypotension would be under general anesthesia (except during methods based upon conduction anesthesia not supplemented by basal narcosis) under these conditions, *cerebral metabolism is reduced almost 30% to 40% and differences between the available oxygen supply and the cerebral oxygen demand are less likely to occur than in the conscious volunteer*

Effects on the Liver The anatomy and physiology of the hepatic blood supply are such that a higher critical level of hypotension exists for the liver than for most other tissues The blood supply to the liver is from two sources, the hepatic artery and the portal vein normally the former supplies about 20% of the total hepatic inflow at a relatively high oxygen tension (95% saturation) whereas the portal vein supplies the remaining 80% of the blood at a much lower (74%) degree of oxygen saturation When the circulation is slowed as during sympathetic paralysis and hypotension a point may be reached at which such a high degree of oxygen desaturation occurs in the intestinal capillaries that the oxygen tension of the blood in the portal vein is nearly as low as that of the blood in the hepatic vein (304) *The liver will then depend entirely upon the hepatic artery for its oxygen supply If the flow through the hepatic artery is reduced it may become inadequate for the total needs of the liver and local tissue anoxia will occur This in fact occurs during hypotension The liver*

which has been subjected to anoxia becomes unable to form urea from ammonium salts or amino acids (281), loses the power to inactivate the vasodepressor material of Shorr (447, 448) and the deterioration of hepatic function is progressive until a point of irreversible change is reached (284)

This is the sequence of events that may occur under controlled hypotension. The estimated hepatic blood flow was decreased 33% from the control value during high spinal analgesia (297) and the hepatic vein oxygen content decreased on average 30% (338). Local anoxia in the liver consequent to the reduced blood flow may then permit the action of V D M: it has been observed during a series of laparotomies conducted under epidural blocks which were sufficiently extensive to cause the blood pressure to fall to 45 to 60 mm Hg that the livers were dark and cyanotic in appearance and turgid and rubbery in consistency; these changes were reversed when the blood pressure was raised by the administration of methedrine or nor adrenaline (64).

Occult liver damage may result from this temporary hypoxia (5). Patients with normal pre-operative liver functions showed abnormal bromsulphalein retention for three to seven days after high spinal analgesia (297) and pathological changes have been shown to occur in the livers of experimental animals after hypotension induced with methonium compounds (432). Indeed hepatic damage may not necessarily be occult: for one fatal case of liver necrosis has been attributed to the use of these techniques (445).

Effects on the kidneys Any consideration of the effects of controlled hypotension on the kidneys must distinguish clearly between the suppression of urine formation during the period of hypotension and the depression of

renal function or the demonstration of morphological changes in the kidney following the restoration of normotension

The formation of urine is by a combination of glomerular filtration and tubular reabsorption. When the blood pressure is lowered to 70 mm Hg or below in the normal human being the hydrostatic pressure in the glomeruli becomes insufficient to maintain filtration and urine formation may be expected to cease therefore when the arterial pressure reaches such a level during controlled hypotension. Both empirical clinical experience and experimental evidence substantiate this hypothesis. During spinal analgesia in the surgical patient renal function is reduced consistently during periods of hypotension (21 297). The same is true when the hypotension has been induced by ganglionic blockade the lowered systemic arterial pressure in both instances causing a reduction of glomerular filtration effective renal blood flow, and urine formation (253 306 308 321 323 475). The extent of these reductions is dependent upon the intensity of the hypotension a moderate fall in blood pressure will produce a decrease in the rate of glomerular filtration for a limited period of time but after an hour or more there will be a rise in the filtration rate despite the continued presence of hypotension (332 335 337) when the decrease in arterial pressure is profound however the inulin clearance rate is at a consistently low level in both man and the experimental animal (317 432).

The suppression of urine formation during the period of hypotension is therefore to be expected if the arterial pressure has been lowered to a significant degree but this fact cannot be construed to mean that renal damage has occurred or will occur. In fact it has been argued that even at blood pressures lower than 70 mm Hg the continued viability of the renal epithelium is not endangered if effective ventilation with oxygen is maintained and the

arteriolar bed is dilated by sympathetic blockade. On the other hand hexamethonium induced hypotension has been shown to cause proteinuria, and the presence of casts and red cells in the urine post-operatively (129) and these findings may (101) be indicative of renal damage (288). At any rate it appears certain that renal damage does in fact occur in view of the high incidence of renal complications and their occasional progress to fatal terminations following the use of the techniques of 'controlled hypotension' (see Chapter V).

The Use of Vasopressor Agents There have been marked differences of opinion regarding the advisability of employing vasopressor drugs when it becomes desirable to restore the blood pressure toward normal levels during or at the end of 'controlled hypotension'. One school of thought has maintained that the blood pressure must be raised prior to closure of the wound in order to permit identification and ligation of all bleeding points since otherwise the danger of reactionary hemorrhage will be great when the pressure finally returns to normal and open vessels begin to bleed. On the other hand there is a second school of thought which feels that to use a vasopressor drug at the end of operation is to defeat the purpose of the technique and that to employ one post operatively is to court disaster (440). The reasoning has been as follows: during hypotension produced by arteriolar dilatation (i.e., sympathetic blockade) clots form in the dilated vessels and when the vessels regain their tone following operation they contract down on these formed clots effectively sealing the vessels; however if a vasopressor agent is administered or the blood pressure rises rapidly from any other cause these clots will be expelled and bleeding will ensue. There is yet a third group of workers who do not employ vasopressor drugs ordinarily but will do so when the occasion warrants their use (Table XI).

Quite aside from the possibilities of reactionary hemor

rhage, however, the use of vasopressor drugs also must be considered in the light of recent knowledge of their actions during both anesthesia and hypotension. Measurements of the contractile force of the heart have been made with

TABLE VI
THE USE OF VASOPRESSOR AGENTS IN
CONTROLLED HYPOTENSION

	<i>Anaesthetists in Great Britain and Ireland</i>	<i>Anesthesiologists in U S A</i>	<i>Totals</i>
Vasopressors not Used	50	28	78
Vasopressors Used	38	63	101
Vasopressors Sometimes Used	90	45	135

strain gauge arches in intact dogs subjected to hexamethonium induced hypotension and indicate that conditions of hypotension associated with a weakened myocardium should dictate the use of such pressor amines as 1 arterenol, methamphetamine or ephedrine, since these drugs will produce an increase in heart force in addition to the well known pressor responses (88). It is also important to remember before employing any vasopressor amines that some of these drugs may precipitate sinister cardiac arrhythmias when administered during anesthesia produced by chloroform, cyclopropane, or trichlorethylene and even in the absence of those anesthetic agents vasopressor drugs may initiate dangerous arrhythmias if the tissues contain an increased concentration of carbon dioxide (101). Carbon dioxide retention to this extent occurs readily of course as a result of the respiratory depression following the induction of high spinal analgesia or the administration of methonium compounds in the techniques of controlled hypotension.

X

Fatal and Non-Fatal Complications

THE PHYSIOLOGICAL derangements inherent in controlled hypotension have resulted in the occurrence of certain fatal and non fatal complications and it has been on the subject of these disasters that the great controversy over these techniques has arisen. As Armstrong Davison has put it, never since the days of the heated controversy about chloroform which divided the anaesthetists of the United Kingdom has there been a matter which has so raised Emotion's end of the see saw with a corresponding depression of the end on which sits Reason. It is particularly striking that the adherents of hypotension seem able to discount adverse effects while they presumably are equally surprised that their opponents see lurking dangers in every shadow (101)

Complications The complication rate following the use of controlled hypotension is amazingly high. The results of questionnaires sent to the members of the Association of Anaesthetists of Great Britain and Ireland revealed this complication rate to be one in every 32 cases the similar questionnaires sent to the Diplomates of the American Board of Anesthesiology reported a complication rate of one in 27 cases and since the total number of complications in the 27,930 cases contained in both series was 908 the over all complication rate was one in every 31 cases (Table XII)

The most common pertinent complication was *reaction*

ary hemorrhage, which occurred in 213 cases and thus constituted 27% of the total number of complications. A very great deal of nonsense has been written on the question

TABLE XII
COMPLICATIONS IN 27,930 CASES OF
CONTROLLED HYPOTENSION

Complication	Great Britain and Ireland 21,125 Cases	U S A 6,805 Cases	Grand Total 27,930 Cases
Anuria	54	8	62
Oliguria	30	24	54
Cerebral Thrombosis	20	9	29
Renal Thrombosis	2	1	3
Coronary Thrombosis	15	3	18
Cardiac Arrest	7	15	22
Cardiovascular Collapse	11	25	36
Reactionary Hemorrhage	164	79	243
Persistent Hypotension	3	70	73
Delayed Awakening	193	3	196
Blurred Vision Post operatively	126	1	127
Others	28	17	45
Totals	653	255	908
Complication Rate	1 in 32	1 in 27	1 in 31

(Reprinted from Little D M Jr Induced hypotension in anesthesia and surgery *Anesthesiology* 16:320 1955 with the kind permission of the publisher)

of reactionary hemorrhage the tendency being to admit the theoretical possibility but to disclaim any knowledge of its actual occurrence. It is worth pointing out that in at least eight of these 243 instances reactionary hemorrhage was not just trivial and annoying but sufficiently severe to cause death (see Table XIV). It is unfortunate that there is no data available to indicate whether it is the use or the avoidance of use of vasopressor agents at the end of surgery that is most likely to be correlated with the occurrence of reactionary hemorrhage (see Chapter IX).

The second most common complication has been a pro

longed and *delayed awakening time* after the operation and anesthesia have been terminated. This phenomenon occurred on 196 occasions and was responsible for 22% of the total number of complications. This slower recovery following controlled hypotension has been attributed in the past to either the synergistic effect which ganglionic blocking agents appear to have with general anesthetic drugs or the lowered metabolic rate induced by these drugs and responsible for their slower elimination from the body (371). Recent studies suggest that there may be a more important cause in the course of observations on electroencephalographic tracings during induced hypotension (65), it was noted that while normal subjects recovered rapidly and completely when the blood pressure rose above 50 mm Hg an arteriosclerotic volunteer showed a delayed return to consciousness and normal cortical rhythm long after the blood pressure had been restored to a much higher level. It is conceivable that the prolonged recovery time seen occasionally in patients who have been subjected to controlled hypotension may be due to cerebral anoxia rather than to the vasodilator drugs themselves.

Blurred vision in the post operative period was the third most common complication and occurred a total of 127 times. It was exhibited exclusively following the use of the methonium preparations and has been attributed to the long lasting action of these drugs upon the eye's power of accommodation. While it was surely an inconvenience, it does not appear to have been a complication that carried with it any prolonged deleterious after effects (73-74).

Fourth in rank as concerns frequency of complications were disturbances involving the kidneys *anuria* occurring 62 times and *oliguria* on 54 occasions. Considering the levels of blood pressure sought in the majority of these 27,930 cases of controlled hypotension (see Chapter IX)

it is rather surprising that there were not more complications involving renal function, but these figures do stress the fact that, of all the vital organs the kidneys are the most liable to damage or dysfunction during induced hypotension

Thrombotic phenomena within essential arterial supplies occurred a total of 50 times *cerebral thrombosis* in 29 patients, *retinal thrombosis* in three patients and *coronary thrombosis* in 18 patients There have also been word of mouth reports of *mesenteric thromboses*, but this complication was not reported on the questionnaires nor has its occurrence been remarked upon in the literature Thrombus formation is an evident danger when the circulation is slowed to the extent that pertains in controlled hypotension'

Cardiac arrest occurred in 22 cases and *cardiovascular collapse* in 36 cases and in roughly 25% of these cases the morbidity progressed to become a mortality (see Table XIV)

Certain other less frequent complications require comment When the blood is circulating through the peripheral vessels with what amounts to a reduced *vis a tergo* as is true during normovolemic hypotension the application of external pressure to the walls of those vessels may constitute a formidable impedance to the flow of blood within the vessel although such pressures would be tolerated well without causing obstruction to circulation under the ordinary circumstances of normotension What has been termed *retractor anemia* is an example in point during the conduct of neurosurgical procedures within the cranium under conditions of controlled hypotension, the retractors employed on the brain substance to provide adequate operative visualization may so restrict the cerebral circulation in the tissues on which they are exerting pressure that cerebral anoxia may result (18) This same situa

tion can occur in other tissues than the brain Pressure necrosis of the laryngeal mucosa has followed the use of an inflated endotracheal cuff in the presence of hypotension In another instance, disastrous aortic femoral thrombosis followed a difficult nephrectomy during which large packs placed in the depths of the wound under consider

TABLE VIII

CORRELATION OF BLOOD PRESSURE LEVELS WITH THE OCCURRENCE OF COMPLICATIONS IN 680 CASES OF "CONTROLLED HYPOTENSION"

Complications	Below 80 Mm 80 Mm Hg or Hg above 3,071 2,664 Cases Cases		<i>p</i>
Anuria	7	0	0.03 > $p > 0.02$? significant
Oliguria	17	5	
Cerebral Thrombosis	9	0	0.02 > $p > 0.01$ significant
Retinal Thrombosis	1	0	
Coronary Thrombosis	2	1	
Cardiac Arrest	14	1	0.01 > $p > 0.001$ highly significant
Cardiovascular Collapse	25	0	0.01 > $p > 0.001$ highly significant
Reactionary Hemorrhage	71	7	0.01 > $p > 0.001$ highly significant
Persistent Hypotension	53	7	0.01 > $p > 0.001$ highly significant

(Reprinted from Hampton L. J. and Little D. M. Jr. *Arch Surg* 1953 with the kind permission of the publisher)

able pressure produced compression of the aorta while at the same time the blood pressure had been deliberately reduced to control bleeding Two cases of unilateral blindness from retinal artery occlusion have occurred apparently by this same mechanism during controlled hypotension pressure on the eyeball by the anesthetic face mask being transmitted through the eye to prevent the flow of blood in the retinal artery (158)

Three major generalities may be made concerning these fatal and non fatal complications First it has become very clear that the extent or degree of hypotension is of paramount importance in respect to the occurrence of morbid

ities and mortalities Hypotension of the order of 80 mm Hg will be tolerated with far fewer complications than will hypotension below that level of blood pressure (Table XIII) Second it is evident that the duration of hypotension is also of significance in respect to the causation of complications 1 1/4 hours has been regarded as the upper limit in terms of the length of time that hypotension may be maintained although it is true that induced hypotension has been continued for far longer periods than this without the occurrence of obvious sequelae There is always the possibility, however, that irreversible shock may follow the use of these techniques and that a method conceived and begun as controlled hypotension may become quite literally uncontrolled hypotension (385) Third experimental studies have suggested that, while the extent and duration of hypotension are of obvious importance the abruptness of onset of hypotension may be a considerable factor in producing complications (425) The maintenance of the integrity of the cardiovascular system is a delicate and finely adjusted mechanism, and an abruptly induced, acute hypotension appears to be more dangerous than a hypotension of equal proportions that is brought on more slowly

Mortalities The deaths which have resulted from the use of the techniques of controlled hypotension have, in general followed the same etiological classifications as the complications and have represented most severe and fatal progressions of those complications In response to questionnaires on the subject British anesthetists reported 46 deaths in 21 125 cases or a mortality rate of one in 459 their American colleagues reported 50 deaths in 6 805 cases or a mortality rate of one in 136 cases and the total number of deaths reported in response to both questionnaires was 96 in 27 930 cases for an over all mortality rate

of one in 291 cases (Table XIV) The great disparity between the mortality rate in the British Isles and that pertaining on the American continent is of interest it may

TABLE XIV
MORTALITIES IN 27,930 CASES OF
"CONTROLLED HYPOTENSION"

<i>Cause of Death</i>	<i>Great Britain and Ireland 21 125 Cases</i>	<i>U S A 6,805 Cases</i>	<i>Grand Total 27,930 Cases</i>
Renal Failure and Anuria	8	1	9
Cerebral Thrombosis or Anoxia	11	7	18
Coronary Thrombosis	7	1	8
Cardiac Arrest or Failure	6	3	9
Cardiovascular Collapse	1	4	5
Reactionary Hemorrhage	2	6	8
Following High Spinal	2	3	5
Over heparinization (Arteriotomy)	1		1
Arterial Air Embolism		1	1
Aortic Femoral Thrombosis		1	1
Pulmonary Infarcts and Edema		1	1
Persistent Hypotension		1	1
Neurosurgical Operative Trauma		14	14
Miscellaneous (Probably not Related)	8	7	15
Totals	46	50	96
Mortality Rate	1 in 459	1 in 136	1 in 291

(Reprinted from Little H M Jr Induced hypotension in anesthesia and surgery *Anesthesiology* 16 320 1955 with the kind permission of the publisher)

indicate that American anesthesiologists are far less skillful than their British counterparts or it may indicate that the patients subjected to controlled hypotension on this continent have been selected for indications that have been so urgent that the calculated risks involved are reflected in the mortality rate

Renal failure and anuria were the cause of death on nine occasions so that in almost 15% of the cases in which anuria was present as a complication a fatality ensued The

following is a typical case of a mortality resulting from anuria, and also serves to emphasize the dangerous results of hemorrhage during 'controlled hypotension

One death must be attributed to mismanagement of the technique. A woman aged 65 years having a combined synchronous abdomino-perineal resection was given 50 mgm hexamethonium bromide. Delay occurred in positioning her on the table and when the abdomen was opened although the blood pressure was at a satisfactory level of 80 mm Hg and there was no bleeding from the abdominal incision the veins of the pelvis were enormously dilated and extensive hemorrhage occurred. Blood was replaced four pints being given but return to normal blood pressure was extremely slow. Methedrine was given in repeated doses of 10 mgm and an adrenaline drip was started which kept her blood pressure at 100 mm Hg. This was continued intermittently for three days. She had complete anuria and died at the end of the third day. (Reported on questionnaire sent to the members of the Association of Anaesthetists of Great Britain and Ireland)

Cerebral thrombosis and anoxia accounted for the largest number of deaths 18, and the following case history is illustrative of this sequelae which caused almost 20% of the reported deaths

A fit old man with a carcinoma of left pinna for excision with block dissection glands of neck. Skin flap from occiput and graft from anterior abdominal wall.

General condition B P 160/85. Electrocardiogram showed slight left ventricular preponderance.

Anaesthetic Omnipon 1/6 grain scopolamine 1/300 grain Thiopentone 0.5 gm curare 15 mgm pethidine 50 mgm oxygen cocaine spray cuffed tube Nitrous oxide-oxygen Respiration assisted at first Feet down tilt 20 degrees Pentamethonium 50 mgm Fifty minute oper

ation during which left jugular vein tied Atropine 1/100 grain Neostigmine 2.5 mgm at end Electrocardiograms throughout hypotensive phase showed no change Consciousness did not return for 4½ hours post-op Patient then quite normal talking taking fluid by mouth moving all limbs B P 120/80 Next morning 24 hours post-operatively found to be semi-conscious speech slurred left facial palsy left arm flaccid plantars extensor Died 3rd post-operative day

P M Very marked atheroma all parts of aorta Left internal carotid commencement completely occluded by clot and atheroma reduced lumen above clot to pin point Right internal carotid similar plaque of atheroma which had reduced lumen considerably—no clot Right middle cerebral artery recent thrombosis fairly near its origin Recent infarct on right parietal lobe Cerebral circulation was obviously impaired depending mainly on vertebral supply (84)

Thrombotic mechanisms were the major factor in causing death on the eight occasions in which they involved the *coronary arteries*, as in the next example

A fat woman with carcinoma of breast for radical mastectomy

General condition Good B P 172/98

Anaesthetic Omnipon 1/6 grain scopolamine 1/300 grain Thiopentone 0.5 gm Nitrous oxide-oxygen trichlorethylene (does not say whether intubated) Pethidine 50 mgm B P 130/84 Hexamethonium iodide 20 mgm Taken into theatre Positioned 10 degrees feet down Pethidine 25 mgm B P 98/70 Operation begun Hexamethonium iodide 10 mgm B P 80/58 Remained steady about this level for 40 minutes when it rose to 90/68 Pulse steady 100-108 Colour good peripheral circulation satisfactory Further pethidine 25 mgm during this time Just as removal of breast being completed (46 minutes after starting) peripheral circulation noted un

satisfactory and B P 50/40 Colour of face grey blue Lungs inflated with oxygen Table 10 degrees head down Methedrine 10 mgm i v Pulse imperceptible Fast plasma drip Further methedrine carbon dioxide 5 per cent oxygen 95 per cent inflation Operation ended at 64 minutes Condition much improved Colour and breathing satisfactory Pulse 120 B P 92/70 Just under one pint of plasma was given during operation Kept in theatre 10 minutes Then as her condition appeared satisfactory patient was transferred on a 10 degree head down trolley to bed which had been blocked Nasal oxygen started Suddenly died 20 minutes post-op

P M Both lungs grossly oedematous Air passages not obstructed Small area of infarction in left ventricle Anterior descending branch left coronary artery almost entirely blocked by atheroma (84)

Cardiac arrest or failure was the cause of nine of the 96 mortalities reported as in this instance

Thirty six year old white male undergoing subtotal thyroidectomy for toxic goiter Hexamethonium employed at the surgeon's request Some signs of toxicity still present in spite of iodine and propylthiouracil treatment Anesthesia induced with pentothal continued with endotracheal cyclopropane and ether C6 given with apparently satisfactory results Thirty minutes later blood pressure fell further and cardiac arrest occurred Cardiac massage performed and later defibrillation Cardiac action was reinstated but the patient failed to recover consciousness and died four hours later Cause of death not definitely determined but presumed to be cerebral and myocardial anoxia (Reported on questionnaire sent to Diplomates of the American Board of Anesthesiology)

Cardiovascular collapse was responsible for death in five patients, and may be exemplified in the following case record

A very old man for emergency excision of carcinoma of larynx

General condition Very poor risk Deep x ray therapy regarded as no avail B P 110/100 Some cyanosis due to obstruction Accessory muscles in use to some extent

Anaesthetic Atropine 1/75 grain Tracheotomy using 1 per cent procaine local Some degree of collapse occurred on inserting cuffed tube Says thiopentone and gallamine given but does not state definitely when Any way lungs inflated with oxygen Recovery in a minute or two Nitrous oxide-oxygen started Blood drip set up Blood pressure then 95/70 Considerable hemorrhage during tracheotomy so decided to use controlled hypotension Hexamethonium iodide 30 mgm Blood pressure fell to 50/40 Slope reduced from 25 degrees to 15 degrees feet down tilt Operation started and proceeded rapidly with negligible bleeding After 40 minutes on lifting larynx to cut through lower part prior to removal patient coughed and respirations became rapid Thiopentone 0.1 Gm and gallamine 8 mgm given but response inadequate Thiopentone 0.05 Gm and gallamine 4 mgm Response satisfactory Few minutes afterwards B P fell from 60/45 to 30/? and within few minutes patient died

P M Small amount of blood in bronchi, and it was thought that this entered during tracheotomy Heart and lungs otherwise in good condition No sign of coronary disease or thrombosis (84)

Reactionary hemorrhage, the most frequent complication also proved to be a major factor in the production of mortality eight deaths occurred for this reason as in the following instance

Middle aged healthy male—hemipelvectomy—ether anesthesia—hexamethonium 225 mgm—head-down position—B P 70/80 mm Hg systolic during greater part of operation—blood loss (measured) about 1000 cc.—re

placed during operation—B P returned to normal at end of operation with neosynephrine drip—patient apparently in good condition when he left O R

After return to ward B P and pulse disappeared A continuous drip of noradrenaline required to maintain circulation until time of death 48 hours later On the night of operation blood volume studies showed a marked deficit of red cells Hemorrhage was suspected and packed red cells were given since plasma volume was relatively good This was in addition to several transfusions given during the immediate post-operative hours Although no abnormal bleeding could be detected at time of blood volume studies on morning of first post operative day blood had distended the abdomen penis and scrotum In spite of blood replacement and noradrenaline drip, his condition steadily deteriorated and he died on the morning of the second post-operative day

Autopsy revealed massive hemorrhage into the peritoneal cavity and external genitalia as well as the operative site No large vessels could be discovered as the cause of bleeding It appeared to have been due to a generalized capillary ooze In addition there were pulmonary edema and acute passive congestion of the lungs central lobular ischemic necrosis of the liver hemorrhagic infarction of the spleen and kidney and hemoglobinuric nephrosis secondary to shock The question arose as to whether these changes were due to C6 or could in part be attributed to prolonged vasoconstriction (Reported on questionnaire sent to Diplomates of the American Board of Anesthesiology)

There were five patients who died following the administration of high spinal anesthesia, as in the following case history

A healthy man with a prolapsed lumbar disc

General condition : Operation five years ago for disc
Good result until seven weeks ago when twisting spine

caused severe recurrence II P 110/70 Apical systolic murmur

Anaesthetic Morphine 1/1 grain scopolamine 1/100 grain Thiopentone 1 Gm No apnea Intubation easy Put into face-down jack knife position Maintenance with nitrous oxide-oxygen trichlorethylene Spinal with heavy nupercaine 1.8 cc/m Head-down tilt increased Blood pressure fell to unreadable levels Respiration rate fell to 8/min Methedrine given 15 Apnoea Inflation with oxygen turned on back heart massage intra aortic saline, more methedrine Only momentary response

P M nothing significant (81)

Finally there were 14 deaths in which the *neurosurgical operative trauma* was considered the major cause of demise or at least could not be excluded as a possible factor of major significance To quote one of the replies to the questionnaires

Two patients have died 48 hours after operation and one patient has died 72 hours after operation In these neurosurgical patients it has been impossible to determine whether death was due to pre existing disease the sequelae of operative trauma or the induced hypotension

This is perhaps an appropriate note upon which to end a discussion of complications and mortalities for there is reason to question the validity of ascribing to induced hypotension the entire burden of guilt in the causation of these complications and deaths herein described It is true that in many instances the techniques of controlled hypotension would appear to be implicated but until there are comparable series of cases run with and without the use of these techniques it is unfair to attribute all deaths occurring in surgical patients in whom hypotension has been induced as being the direct result of that hypotension

XI

Contraindications

INSIGHT into the contraindications to the employment of controlled hypotension follows most naturally upon knowledge of the physiological aberrations produced by these techniques and the morbidities and mortalities consequent to their use. The major concern, not unreasonably, has been expressed over the state of the cardiovascular system particularly the presence of disease in the various branches of the vascular tree, and over the status of the blood volume and the oxygen carrying capacity of the blood within those vessels. In the questionnaires sent to the members of the Association of Anaesthetists of Great Britain and Ireland and to the Diplomates of the American Board of Anesthesiology each physician was requested to designate his considered contraindications to the use of these techniques (Table XV).

The presence of *arteriosclerosis*, known *coronary disease* or previous *cerebrovascular disease* must be considered absolute contraindications to the employment of controlled hypotension since there is great danger that the flow of blood through vessels which are already partially occluded may cease altogether under conditions of hypotension. The risk of coronary thrombosis or occlusion is very real during hypotension, even in the absence of known coronary insufficiency (203) and in spite of the hypothesis that vasodilatation and lowered peripheral resistance will reduce the work load of the heart (175). Furthermore since the circulation time is increased almost two fold during induced hypotension thrombus formation

is very likely and especially so in the presence of narrowed vessels. The fact that cerebral blood flow, for example, may be maintained near normal in moderate degrees of hypotension by a decrease in cerebral vascular resistance does

TABLE XV

CONTRAINDICATIONS TO CONTROLLED HYPOTENSION

	Anaesthetists in Great Britain and Ireland	Anesthesiologists in U S A	Totals
Cardiovascular Disease			
In General	48	19	67
Coronary Disease	62	57	119
Arteriosclerosis	51	30	81
Cardiac Disease	46	45	91
Cerebrovascular Disease	32	22	54
Hypertension (Severe)	43	24	67
Hepatic Disease	8	14	22
Renal Disease	48	40	88
Extremes of Age	82	31	113
Disturbances of Blood Volume			
Anemia	18	18	36
Hemorrhage	10	7	17
Shock	40	12	52
Low Blood Volume	30	19	49
Polycythemia	3	4	7
Uncomplicated Elective Surgery	54	17	71
Others	58	59	117

(Reprinted from Little H M Jr. Induced hypotension in anesthesia and surgery *Anesthesiology* 16 320 1955 with the kind permission of the publisher)

not necessarily mean that there is less likelihood of cerebral thrombosis for if cerebral vascular resistance is reduced by dilatation of the cerebral vessels which is the mechanism that must be assumed unless blood viscosity or intracranial tension are considered to vary the slowing of the linear flow rate and the prolongation of the local circulation time will enhance the possibility of the development of intravascular thrombosis (103)

Cardiac disease in general and particularly cardiac failure have been considered contraindications to the use of

these techniques and in the present state of knowledge this is undoubtedly the correct view. It is worth noting in passing, however, that the cautious use of ganglionic blocking agents in the therapy of cardiac failure is now under trial, based on the same rationale as the use of phlebotomy or spinal block in the past, and hypotensive techniques may well play an active part in the anesthesia for certain cardiac patients in the future.

Severe and malignant *hypertension* must be considered an absolute contraindication to controlled hypotension, but only when excessively severe. In all other forms, hypertension is only a relative contraindication, if the exaggerated effects of hypotension upon cerebral blood flow, renal blood flow, and other parts of the cardiovascular mechanism in the hypertensive subject are kept in mind.

The presence of *hepatic disease* must certainly constitute an absolute contraindication, as must the presence of *renal disease*. Liver failure is liable to follow extreme hypotension due to the peculiarities of the hepatic blood supply, and a case of acute hepatic necrosis has already been reported following the use of methonium induced hypotension (445). Tests of renal function during and after controlled hypotension demonstrate quite clearly the adverse effects that these techniques may have on the kidneys when the latter are in a diseased state. There is a further consideration in connection with kidney function which was mentioned previously and may be extremely important: both the methonium compounds and arfonad depend at least in part upon the kidneys for their excretion. In the presence of profound hypotension, glomerular filtration is reduced to such an extent that urinary excretion of these drugs does not occur; a vicious cycle is then set up in which failure of the kidneys to excrete these ganglionic blocking agents produces a continuance of their hypotensive effects. As would be expected, this is particu-

larly prone to occur when the kidneys themselves are diseased

A further absolute contraindication to the use of the techniques of 'controlled hypotension' is a disturbance of blood volume and oxygen carrying capacity of the blood such as *anemia*, *hemorrhage*, and *low blood volume*. *Shock* and *hypotension* are usually part of the clinical picture of such disturbances of blood volume and are, of course contraindications in themselves. In Addison's Disease, hypotension exists already, and although the blood volume may be relatively normal further reduction of the blood pressure must be considered unsafe. Extreme increase in blood volume is also a contraindication because of the markedly slowed circulation the tendency toward thrombotic episodes in patients with *polycythemia vera* must not be aggravated through further slowing of the circulation by the induction of hypotension.

There is another absolute contraindication of importance. If the operating room team, the personnel responsible for post operative care or the facilities for either operative or post operative care are in any way deficient controlled hypotension should not be employed. These are not techniques for the occasional operator or anesthetist they are not techniques for inexperienced or slow surgeons and they are not techniques which can be employed safely when routine post operative care is the only such care available.

The extremes of age have often been considered as contraindications but it would appear that these are only relative contraindications. Patients under sixteen years of age are often resistant to ganglionic blocking drugs particularly the methonium compounds and it is therefore occasionally difficult to induce a satisfactory hypotension in such patients. This would appear to constitute a failure of the technique rather than a contraindication to its

use, excepting that this failure can on occasion be dangerous of itself for if the ganglionic blockade produces partial vasodilatation, without a marked fall in blood pressure, operative bleeding may be actually increased (120) The use of these techniques in patients over 60 years of age has also been considered a contraindication, because of the possibility of unsuspected and latent cardiovascular disease (203) However the absolute chronological age of the patient is far less important than the physiological age of his or her arterial tree

Diabetes Mellitus is also a relative contraindication, at least to the use of the methonium compounds for there is a tendency for the blood sugar to fall during the administration of these drugs The hypoglycemia which occurs as a result may become very profound without being recognized since the usual symptomatology of sweating, palpitation, restlessness and anxiety, are due to sympathetic and adrenal discharge and hence do not occur in the presence of ganglionic blockade The accentuation of the effects of the administration of insulin in the diabetic patient by the methonium compounds together with the masking of these effects by general anesthesia, can result in a very dangerous situation (12 176)

Without question however, the most absolute of all contraindications to the use of the techniques of 'controlled hypotension' must be the *absence of a rational and urgent indication* There can be no justification for the use of induced hypotension merely for convenience during uncomplicated and elective surgery Controlled hypotension is not a luxury during surgery it is an occasional necessity and must be employed only under conditions of the most compelling need There are grave dangers associated with its use but there are also on occasion, great advantages to be attained from that use The method must therefore be considered contraindicated in those

situations in which the advantages do not outweigh the risks involved

These contraindications just listed are of immense importance and failure to respect them can lead to disaster. They do not represent merely a list of pathological entities which might conceivably be of theoretical danger; they are conditions which bitter empirical experience has taught are incompatible with the successful conduct of 'controlled hypotension'.

XII

Indications

THE INDICATIONS for which controlled hypotension is employed are the crux of the problem concerning these techniques. On what basis does one seek to induce hypotension and when does one fight to maintain a normal arterial pressure? There is no glib answer to this riddle unless it were to be Sir John Gillies' statement to the effect that there is only one true indication for induced hypotension—that situation in which its advantages will be of certain benefit to the patient and likely to outweigh the assumed risks.

In any discussion of the indications for the use of controlled hypotension, therefore, it is imperative to make a clear distinction between the specific operative procedures for which these techniques have been employed in the past and the clinical situations for which their use is justifiably indicated. Controlled hypotension has been advocated for a large variety of operative procedures including such unlikely choices as tonsillectomy, dental extractions and caesarean sections. Most commonly, however, it has been employed during extensive surgical interventions in which there could be expected to be a large amount of bleeding, sudden or profuse hemorrhage or greatly increased tension within vessels or organs.

The categorical listing of these procedures is a long one (Table XVI): *neurosurgical procedures* (18, 26, 27, 80, 187, 188, 189, 207, 228, 318, 322, 375, 376, 378, 381, 394, 451, 474, 491, 492, 501, 515), particularly the removal of highly vascular tumors; (4) the ligation of various types of

intracranial aneurysms (377, 395, 466, 508), and the removal of meningiomas (151), *head and neck surgery*, such as the Lempert fenestration operation (135, 168 200 224, 347, 494) ophthalmic surgery (399) in the presence of glau

TABLE XVI

TYPES OF OPERATIVE PROCEDURES FOR WHICH
CONTROLLED HYPOTENSION HAS BEEN EMPLOYED

Neurosurgery	Vascular Brain Tumor Congenital Aneurysm Meningioma
Head and Neck Surgery	Lempert Fenestration Operation Ophthalmic Surgery Radical Neck Dissection Thyroidectomy Laryngectomy
Surgery of Chest Wall	Radical Mastectomy Thoracoplasty
Intra thoracic Surgery	Pneumonectomy Lobectomy Decortication Esophago gastrectomy
Cardiovascular Surgery	Coarctation of Aorta Patent Ductus Arteriosus Aortic Aneurysm
Abdominal Surgery	Radical Abdominal Operation Abdomino perineal Resection Gas trectomy Porta caval Shunt
Pelvic Surgery	Pelvic Evisceration Radical Hysterectomy Pelvic Node Dissection
Orthopedic Surgery	Spinal Fusion Laminectomy Open Hip Operation Bone Graft
Genito urinary Surgery	Prostatectomy Transurethral Resection Nephrectomy Adrenalectomy Vesico vaginal Fistula Vaginal Plastic Operation
Plastic Surgery	Skin Graft Skin Flap

coma (73 211), a number of otorhinolaryngological procedures (155 168, 282, 355 506) thyroidectomy (231 398) radical neck dissection (223 231) and laryngectomy *surgery of the chest wall*, specifically radical mastectomy (231) and thoracoplasty (398) *intra thoracic surgery* (105 291 422) such as pneumonectomy (79 96) lobectomy decortication and esophago-gastrectomy (67) *cardiovascular surgery* (110 210) particularly that involving large vessels (166) such as coarctation of the aorta patent ductus arteriosus and aortic aneurysms *abdominal surgery* (58, 59 60 111) including radical abdominal operations,

gastrectomy abdomino perineal resection thoraco lumbar sympathectomy (175 240) and porta caval shunt *pelvic surgery* (58 59 60 270) particularly *pelvic evisceration* radical hysterectomy and pelvic node dissection *ortho* *pedic surgery* (462) such as spinal fusion, laminectomy (382) open hip operation (242) and bone grafts *urologi* *cal surgery* (497) including *adrenalectomy* *nephrectomy* *prostatectomy* (50 85 86 87 325 393) *vesico vaginal* *fistula* and *vaginal plastic operation* and *plastic surgery* (43 373) particularly *skin grafts* and *skin flaps* It must be emphasized that long as this list is it is by no means complete for there is scarcely an operative procedure to which the techniques of controlled hypotension have not been applied

Yet there are no specific operative procedures for which the use of controlled hypotension is indicated and an enumeration of the surgical interventions for which it has been employed must never be confused with the indications for its use There are as previously stated a number of well defined clinical situations during operations in which the use of these techniques may become mandatory and without which the operations would be totally unfeasible immensely more dangerous or more likely to result in unsuccessful outcomes these are the indications for the use of controlled hypotension (Table XVII)

Surgical Indications Foremost among the indications for the use of controlled hypotension must be the *control* of gross hemorrhage during operative procedures for such lesions as intracranial aneurysms vascular brain tumors and definitive interventions for the excision of other highly vascular lesions which are capable of sudden and profuse hemorrhage An indication of less dramatic but equally important significance has been the *control* of *extensive capillary ooze* during operations of considerable magnitude such as abdominal or pelvic *eviscerations* or

radical node dissection The amount of blood lost need not necessarily be great to require the use of induced hypotension the Lempert fenestration operation for example, entails very little actual blood loss yet the techniques of

TABLE XVII
INDICATIONS FOR CONTROLLED HYPOTENSION

Surgical

- Control of gross hemorrhage—i.e. intracranial aneurysm
- Control of extensive ooze—i.e. abdominal evisceration
- Bloodless field for visibility—i.e. fenestration
- Reduction of organ tension—i.e. craniotomy
- Reduction of intravascular tension—i.e. cardiac and great vessel surgery
- Control of hypertensive crises—i.e. pheochromocytoma
- Avoidance of transfusions—i.e. compatible blood unavailable history of transfusion reactions

Anesthetic

- Obtundation of noxious stimuli by ganglionic blockade
- Artificial hibernation

Medical

- Treatment of pulmonary edema in hypertensive crises
 - Therapy in pulmonary embolism
 - Pain relief in herpes zoster
 - Relief of venous congestion in cardiac failure
 - Treatment of severe hemorrhages
-

controlled hypotension may be indicated, for the purpose of increasing *visibility in the operative site*, by the presence of even a single drop of blood which can obscure the entire operative field during the delicate maneuver of shaping the fenestra and thus negate a successful surgical result

The indications for the usage of controlled hypotension must also include those situations in which although bleeding may be expected to be minimal, the tensions within the vessels or organs themselves will be of such magnitude as to endanger the life of the patient or to render improbable a satisfactory completion of the surgical procedure During the course of craniotomies in patients with hugely swollen and edematous brains these

techniques can be employed to *reduce organ tension* and so to facilitate surgical exposure, which might otherwise be totally unsatisfactory, and the same may be true of operations upon the glaucomatous patient, in whom the reduction of arterial pressure leads to a secondary reduction of pressure within the eyeball and permits the performance of operations that would under other circumstances be highly dangerous. A similar situation pertains during certain instances of cardiac and great vessel surgery, in which, by the *reduction of intravascular tension*, it may become possible to place clamps upon tense and bulging vessels, which would tear and precipitate massive hemorrhage were the tensions within the lumens not reduced by controlled hypotension. The reduction of vascular tension may be indicated also in certain rarer instances of surgery during which *hypertensive crises* may threaten to explode quite literally into hemorrhagic accidents a situation encountered in such surgery as the removal of pheochromocytoma (83).

There are certain less frequent indications for the use of controlled hypotension during anesthesia and surgery which should be cited. One such indication is the *avoidance of transfusion* emergency surgery of an acute nature may be necessary in the absence of compatible whole blood for transfusion, and the use of these techniques can prevent blood loss to a degree that transfusion may be avoided altogether. A similar indication may arise in the rare patient with a history of multiple transfusion reactions who must undergo extensive surgery the use of controlled hypotension in such a patient may circumvent the danger of a major transfusion incompatibility by decreasing blood loss to such an extent that transfusion is not essential (327). These would be most infrequent indications under modern operating conditions.

Anesthetic Indications One interesting outgrowth of

the use of controlled hypotension in anesthesia and surgery has been *the use of ganglionic blockade to prevent the transmission of noxious impulses*. This is not, of course a true indication for induced hypotension itself, but it has been suggested that the future may prove that this usage of ganglionic blockade is the most important and enduring contribution that the techniques of "controlled hypotension" have made to medicine. The basis of this use of ganglionic blockade to prevent the transmission of noxious impulses harks back to the theory of Anoci association pronounced by George Crile, senior some 40 years ago. Crile, reasoning along physiological lines, argued that not only must the brain be protected against destructive psychic strain by the use of general anesthesia, but that local anesthesia must be employed also to exclude the noxious impulses arising from the site of surgical intervention (90). It is now apparent that many of these noxious impulses are conveyed by the autonomic nervous system and therefore are susceptible to ganglionic blockade (57, 82, 389). The clinical impression has been gained that patients subjected to controlled hypotension develop surgical shock less frequently than do other surgical patients (510), and that obtundation of noxious impulses by ganglionic blockade is feasible (39, 436).

The French have made specific utilization of these ideas in their techniques of artificial hibernation in which induced hypotension is one of the several essential factors in a combination that includes hypothermia, central nervous system depression, muscle relaxation, and that evasive delight *déconnexion* (484).

Medical Indications. Finally and similarly, an increasing number of medical therapies have been based upon the use of the ganglionic blocking agents which have achieved prominence through their employment in the techniques of controlled hypotension, the treatment of

pulmonary edema in hypertensive crises (95, 415 416 417 418, 419) *the therapy of pain and neurogenic shock in pulmonary embolism* (91, 429) *pain relief in herpes zoster* (346), *the potential relief of venous congestion in cardiac failure*, (505) and *the treatment of severe hemorrhages* (109, 199, 280) and *cerebrovascular bleeding* (430) a revolutionary but apparently effective therapy. The further extension of the use of ganglionic blockade in the therapy of a number of other pathological processes seems assured.

XIII

Requirements

THE CONDUCT of a surgical intervention performed under the techniques of controlled hypotension demands meticulous attention to detail and rigid adherence to a set of fundamental tenets. Any lapse in the fulfillment of the requirements is a sure invitation to disaster, for these are techniques which are quite capable of either maiming or killing the patient. This is not to say that they should be forbidden techniques for most assuredly the results of their use can often be useful sometimes beneficial, and on occasions truly life saving but these are services that are achieved at a price and at times the price may be inordinately high. There is a growing tendency throughout medicine but perhaps particularly in anaesthesia and surgery to equate what is new with what is good and to confuse the means with the end—especially where the means are striking and dramatic. It therefore devolves upon those who induce hypotension deliberately in the surgical patient to adopt rigid criteria for the execution of the details of controlled hypotension (Table XVIII) for the usefulness of these techniques is entirely dependent upon their successful performance.

The most unyielding of these requirements without which all the others become meaningless is the absolute necessity for selecting patients to be subjected to these techniques on the basis of the most *urgent and rational indications*. Such indications must be based upon the probable failure either of the patient to survive the operation or of the surgery to be ~~completed successfully~~ without the

use of controlled hypotension." The use of these techniques in the absence of a rigidly-defined indication is at best poor judgment and is at worst closely akin to malpractice.

As a corollary to the requirement that there must be the most scrupulous selection of cases on the basis of indi-

TABLE XVIII
THE REQUIREMENTS OF CONTROLLED HYPOTENSION

-
- | | |
|---|---|
| 1 | A rational and urgent indication |
| 2 | Respect for existing contraindications |
| 3 | Restriction of extent and duration of hypotension |
| 4 | Normal pre-operative blood volume |
| 5 | Drop by drop replacement of blood and fluid loss |
| 6 | Adequate oxygenation |
| 7 | Due consideration to posture |
| 8 | Completely competent anesthesiologist and surgical team |
| 9 | Detailed and meticulous post operative care |
-

cations that are so urgent that operation could not be performed successfully in the absence of the employment of these techniques it is mandatory that any *existing contraindications be respected* in the most assiduous manner. These contraindications must be considered to include disease of various branches of the vascular tree: hepatic and renal disease, and blood volume disturbances of a number of different types (see Chapter XI). A failure to recognize the existence of these pathological entities prior to the induction of hypotension in the surgical patient has been responsible, in the past, for some of the tragic morbidities and mortalities which have followed upon the use of controlled hypotension. The occasional clinical situation does arise in which the need and indication for controlled hypotension is so overwhelming that the physician's judgment dictates that an existing contraindication must be ignored: such situations are indeed rare, however.

When a pregnant indication for the use of controlled hypotension does exist and the decision is made to em-

ploy the technique, it is essential that the *blood pressure be lowered slowly to the maximal level compatible with the control of bleeding, and that it be maintained at this lowered level the shortest time consistent with the adequate accomplishment of the surgical intervention*. It has become very clear that the morbidities and mortalities associated with the use of induced hypotension have been directly related to both the extent and the duration of the reduction of the blood pressure from the pre operative normal. From a purely realistic and practical standpoint, it is almost always possible to restrict the period of hypotension to that part of the operative procedure for which its benefits are most necessary, and advantage should be taken of this fact. Furthermore since the level of the blood pressure which must obtain in order to provide a relatively bloodless field varies tremendously between individual patients the degree of the hypotension should never be greater than that required in the patient under consideration.

These techniques are based upon the production of either a normovolemic or a hypovolemic hypotension and in both instances a point is reached at which the wide spread transgressions of the physiological status quo of the cardiovascular system may progress to irreversible shock. This point may be reached with deceiving rapidity when the blood volume is abnormally low at the time of the induction of hypotension or if it is lowered by hemorrhage during the course of the conduct of controlled hypotension. For this reason it is a requirement that there be an absolutely *normal pre operative total blood volume* and red cell mass. Furthermore it is necessary to recognize that a low pre operative blood volume often cannot be distinguished on purely clinical grounds or even by the usual laboratory investigations so that the selection of patients should be made on the basis of blood volume determinations. This is particularly true of the older

cachectic patient with extensive malignant involvement, who represents a state which has been termed 'chronic shock

By the same token hemorrhage or extensive bleeding which occurs during the course of controlled hypotension also will lead rapidly to the point of irreversible shock, for it has become apparent that patients in whom hypotension has been induced tolerate blood loss poorly. A further requirement therefore must be the *drop by drop replacement of all blood and fluid that is lost* during operation. It should be unnecessary to add that an indwelling cannula into some part of the vascular tree is quite indispensable.

When hypotension has been induced, the oxygen-carrying capacity of the blood must be utilized to the fullest extent if tissue oxygenation and cellular respiration are to be maintained near normal. The accomplishment of this task of *adequate oxygenation* requires a completely patent airway at all times, high flows of oxygen and adequate (preferably spontaneous) ventilation of the lungs.

Due consideration to the posture of the patient is extremely important in controlled hypotension. This matter is of concern not only to promote venous drainage from the wound but also in order to effect the necessary level of hypotension in many instances. At the same time it must be remembered that the position of the patient is often the key to morbidity or mortality in these techniques. Certain postures such as steep Fowler's position, have an undesirable effect upon the blood flow to vital organs particularly the brain. Furthermore shifts of posture in the presence of a dilated vascular bed carry their own inherent risks since a further and more severe hypotension can be so induced with a consequent reduction of venous return to a point incompatible with continued circulatory function.

These heroic maneuvers of controlled hypotension

demand the utmost from those employing the techniques, which means that there must be a *completely competent surgical team*. Not only must the anesthesiologist be highly trained, skilled, and thoroughly conversant with the technique employed, but also the operating surgeon and his assistants must be dextrous, accomplished, and possessed of a complete understanding of what is being undertaken. In perhaps no field of surgical endeavour is there greater need for the utmost in cooperation between all members of the operating team.

Since the dangers to the patient during the immediate post-operative period are at least as great as during the operation and the period of hypotension itself, a final requirement is for *detailed and meticulous post operative care*. Posture, airway, oxygenation, determinations of hematocrit and total blood volume, and constant observation, are all of concern in the post operative period, for the end of operation does not necessarily signify the end of ganglionic blockade nor the attainment of a normotensive state.

In a recent survey studying the effects of anesthesia and surgery in 599 548 surgical patients the anesthetic death rate was found to be one in 1560 patients (36). In 27 930 cases of controlled hypotension, the over all mortality rate has been found to be one in 291 patients. This is indeed a staggering figure. Let there be no doubt that these are gravely dangerous techniques, possessing those inherent capacities for tragedy which always accompany marked deviations from the normal physiological status quo. It would be at least convenient merely to dismiss controlled hypotension as a menace to humanity unfortunately the situation is scarcely that simple. For these are techniques which although potentially lethal are capable of immense benefit for the patient when employed for good and sufficient reason by the knowing, the wary, and the skilled.

XIV

Bibliography*

- 1 ADRIANI J Hypotensive anesthesia *J Am A Nurse Anesthetists* 22 24 1954
- 2 ————— Some recent items of interest to laryngologists concerning anesthesia *Ann Otol Rhin & Laryng* 62 423 1953
- 3 ALEXANDER G L Death after anaesthetic with hypotension *Lancet* 1 299 1953
- 4 ANDERSON S and McKISsock W Controlled hypotension with arfonad in neurosurgery with special reference to vascular lesions *Lancet* 2 754 1953
- 5 ANLYAN W G SHINGLETON W W BENSON W R STEPHEN C R SALEM M and TAYLOR H C A study of liver damage following induced hypotension *Surgery* 36 375 1954
- 6 ANNOTATION Arteriotomy *Brit.M.J* 1 965 1952
- 7 ————— Procaine amide *Brit.M.J* 1 265 1953
- 8 ————— Dangers of induced hypotension *Brit.M.J* 1 324 1954
- 9 ————— Bleeding at operations *Lancet* 1 677 1951
- 10 ————— BP up or BP down? *Lancet* 2 161 1951
- 11 ————— Control of bleeding in cranial operations *Lancet* 2 393 1951
- 12 ————— Diabetes and controlled hypotension *Lancet* 1 404 1954
- 13 APGAR V and PAPPER E M Pheochromocytoma anesthetic management during surgical treatment *A.M.A Arch Surg* 62 634 1951
- 14 ARNOLD P GOETZ R H and ROSENHEIM M L Effect of pentamethonium iodide on the peripheral circulation *Lancet* 2 408 1949
- 15 ARNOLD P and ROSENHEIM M L Effect of pentamethonium iodide on normal and hypertensive persons *Lancet* 2 321 1949
- 16 ARROWOOD J G Differential spinal block With particular reference to hypertensive patients *Proc Roy Soc Med* 43 919 1950
- 17 ————— and SARNOFF S J Differential spinal block V Use in

* This is a compilation of the studies and reports embracing the subject of controlled hypotension. It includes all the works referred to in the text as well as many others which the reader may wish to peruse for additional information.

- the investigation of pain following amputation *Anesthesiology*, 9 614 1948
- 18 ASERMAN E Controlled hypotension in neurosurgery with hexamethonium and procaine amide *Brit.MJ*, 1-961 1953
 - 19 ASQUITH E Controlled hypotension in surgery *Canad. M.A.J.* 67 421 1952
 - 20 ASSALI N S DOUGLASS R A JR and SUYEMOTO R Observations on the hemodynamic properties of a thiophanium derivative RO 2 2222 (arfonad) in human subjects *Circulation* 8 62 1953
 - 21 ASSALI N S KAPLAN S A FOMON S J DOUGLASS R A and TADA Y The effect of high spinal anesthesia on the renal hemodynamics and the excretion of electrolytes during osmotic diuresis in the hypotensive normal pregnant woman *J Clin Investigation* 30 916 1951
 - 22 ASSALI N S and SUYEMOTO R The place of hydrazinophthalazine and thiophanium compounds in the management of hypertensive complications of pregnancy *Am J Obst & Gynec* 64 1021 1952
 - 23 BAARS J H Kunstmatige bloeddrukverlaging bij gynaecologische operaties *Nederl tijdschr verlosk en gynaec* 53 335 1953
 - 24 BACHER K R Pendiomid (Ciba 9295) vor und während der spinalanästhesie und weitere beobachtungen *Zentralbl Chir*, 78 1891 1953
 - 25 ——— Hypotensive periduralanästhesie *Med Klin Berlin* 49 622 1954
 - 26 BALLANTINE R I W Controlled hypotension in neurosurgery *Brit.MJ* 1 1164 1953
 - 27 ——— and JACKSON I Anaesthesia for neurosurgical operations *Anaesthesia* 9 4 1954
 - 28 ——— and BOULTON T B Controlled hypotension with arfonad *Lancet* 1 778 1954
 - 29 BANDIERA C Lipotensione controllata in chirurgia *Rassegna internaz clin e terap* 32 686 1952
 - 30 BANTON A H Methonium compounds in hypertension *Lancet* 1 527 1951
 - 31 BARBIER P Les bases physiologiques de l'emploi des méthoniums en clinique *Semaine méd* 28 639 1952
 - 32 BARLOW R B and ING H R Curare like action of polymethylene bis quaternary ammonium salts *Nature London* 161 718 1948
 - 33 BARNETT A J Comparison of pentamethonium and hexamethonium bromide *Lancet* 1 1415 1951
 - 34 ——— Ocular effects of methonium compounds *Brit J Ophth* 36 593 1952
 - 35 BEACONSFIELD P Dangers de l'emploi de l'hexaméthonium en médecine et en chirurgie *Presse méd* 60 811 1952
 - 36 BEECHER H K and TODD D P A study of the deaths associated

- with anesthesia and surgery Based on a study of 599,548 anesthetics in ten institutions 1948 1952 inclusive *Ann Surg* 140:2 1954
- 37 BEIN H J and MEIER R Zur pharmakologie des NNN N 3 pentamethyl N N' diethyl 3 aza pentan 1,5 diammonium dibromid (Ciba 9290) einer ganglionär hemmenden substanz *Experientia Basel* 6 351 1950
- 38 ————— Pharmakologische untersuchungen über pendiomid eine neuartige substanz mit ganglienblockierender *Schweiz med Wchnschr* 81 446 1951
- 39 ————— Zur frage der schockbekämpfung mit pendiomid *Anaesthesist* 3 25 1954
- 40 BELFRAGE D Death after anaesthetic with hypotension *Lancet* 2 1183 1952
- 41 BENKE A Zur kontrollierten blutdrucksenkung mit depressin und pendiomid *Wien med Wchnschr* 103 187 1953
- 42 BENNETT H S BASSETT D L and BEECHER H K Influence of anesthesia (ether cyclopropane sodium evipal) on the circulation under normal and shock conditions *J Clin Investigation* 23 181 1944
- 43 BENTEL H and PENN J Hypotensive anaesthesia in plastic surgery a report on 500 cases *South African M J* 27 405 1953
- 44 BERNSTEIN A ESSER H and LORENZ W klinisch experimentelle und therapeutische beobachtungen mit pendiomid einer neuen ganglienblockierenden substanz *Schweiz med Wchnschr* 81 452 1951
- 45 BERNSTEIN A and SIEMONS E Der hirnkreislauf bei der gesteuerten experimentellen hypotension (hypotension contrôlée) *Schweiz med Wchnschr* 83 210 1953
- 46 BERTOLLOTTI R Lipotensione controllata in anestesia *Acta anaesth Padova* 3 141 1952
- 47 BIEDERMANN G Prostatachirurgie und willkürliche blutdrucksenkung *Med Klin Berlin* 47 651 1952
- 48 BILSLAND W L Controlled hypotension by arteriotomy in intracranial surgery *Anaesthesia* 6 20 1951
- 49 BIZARD G VANLERINBERGHE J GREGOIRE N ROBELET A and MILBLED G Oxy-métrie au cours de l'hypotension contrôlée par le pentaméthonium *Lille chir* 7 204 1952
- 50 BLAKELEY C G Controlled hypotension *Lancet* 2 198 1952
- 51 BLALOCK A The effects of ether chloroform and ethyl chloride anaesthetics on minute cardiac output and blood pressure an experimental study *Surg Gynec & Obst* 46 72 1928
- 52 BLOCK M A WAKIM K G MANN F C and BENNETT W A Renal lesions and function following prolonged experimental hypotension *Surgery* 32 551 1952

- 53 BLUM R Die prostatektomie in hypotension *Ztschr Urol* 46:761 1953
- 54 BODMAN R I Death after anaesthetic with hypotension *Lancet* 2 1085 1952
- 55 BORRI B L and NATELLIS F Bloodletting during operation to diminish bleeding *Anesth & Analg* 32 6 1953
- 56 BOURNE W LEIGH M D INGLIS A N and HOWELL, G R Spinal anesthesia for thoracic surgery *Anesthesiology* 3:272 1942
- 57 BOYAN C P Ganglionic blockade with methonium compounds *Brit.M.J* 1 728 1953
- 58 ——— Hypotensive anesthesia for radical pelvic and abdominal surgery *A.M.A Arch Surg* 67 803 1953
- 59 ———, and BRUNSWIG A Hypotensive anesthesia in radical pelvic and abdominal surgery *Surgery* 31 829 1952
- 60 ——— ——— ——— Lanesthésie sous hypotension dans la chirurgie radicale des cancers avancés *Lyon chir* 48 37 1953
- 61 BOZZA M Ipotensione controllata mediante un derivato del tiopano esperienza su 55 casi di neurochirurgia *Acta anaesth Padova* 4 389 1953
- 62 BROMAGE P R Vascular hypotension in 107 cases of epidural analgesia *Anaesthesia* 6:26 1951
- 63 ——— Lumbar epidural analgesia for major surgery below the diaphragm A report of 355 cases *Anaesthesia* 7 171 1952
- 64 ——— Effect of induced vascular hypotension on the liver Alterations in appearance and consistence *Lancet* 2 10 1952
- 65 ——— Some electroencephalographic changes associated with induced vascular hypotension *Proc Roy Soc Med* 46 919 1953
- 66 BROWN A S Pre operative exsanguination in intracranial surgery *Anaesthesia* 9 17 1954
- 67 BRUNET D AUBIAC B L'emploi de l'hexaméthonium dans la chirurgie digestive à thorax ouvert *Anesth et analg* 9 400 1952
- 68 BRUNSWIG A Hypotensive anesthesia *Am J Surg* 83 1 1952
- 69 ——— and BOYAN C P Hypotensive anesthesia *West Virginia M J* 49:245 1953
- 70 BULLOUGH J A T piece technique for general inhalational anaesthesia A safe method of reducing operative bleeding *Brit M J* 1 28 1952
- 71 BURT C C and GRAHAM A J P Pentamethonium and hexamethonium iodide in investigation of peripheral vascular disease and hypertension *Brit.M.J* 1 455 1950
- 72 CAMERINI F GUCLIELMI F and PIZZOLI R Studio elettrocardiografico nel corso di interventi operatori in ipotensione controllata *Gior ital anestesiol* 18 503 1952

- 73 CAMERON A J and BURN R A Hexamethonium and glaucoma
Brit J Ophth 36 482 1952
- 74 CAMPBELL A and ROBERTSON I Treatment of severe hypertension
with hexamethonium bromide *Brit M J* 2 804 1950
- 75 ————— Hexamethonium bitartrate *Lancet*, 1797
1951
- 76 CARBONERA D and RESEGOTTI E Lipotensione controllata con ar
fonad *Acta anaesth Padova* 4 377 1953
- 77 CARETTE L Effets rénaux de l'hypotension contrôlée sous anesthésie
générale *Anesth et analg* 11 125 1954
- 78 CARRAVETTA M and RIZZI R Lipotensione nella anestesia chirur
gica *Gior ital anestesiol* 18 445 1952
- 79 CHATEAUREYNAUD J I Méthode pondérale d'évaluation de l'hémor
ragie chirurgicale.—II Utilisation systématique des sels d'hexamé
thonium en chirurgie pulmonaire *Anesth et analg* 9 353 1952
- 80 CHRISTENSEN J C Hipotensión controlada en neurocirugía medi
ante pendiomid *Prensa méd argent* 40 1695 1953
- 81 CHURCHILL DAVIDSON H C WYLIE W D MILES B E and DE
WARDENER H E The effects of adrenaline noradrenaline and
methedrine on the renal circulation during anaesthesia *Lancet*
2 803 1951
- 82 CILIBERTI B J GOLDFEIN J and ROVENSTINE E A Hypertension
during anesthesia in patients with spinal cord injuries *Anesthe
siology* 15 273 1954
- 83 COBB G C HALL J and HALL R A Control of blood pressure in
operation for pheochromocytoma *Lancet* 1 973 1953
- 84 COMMITTEE UPON DEATHS ASSOCIATED WITH ANAESTHESIA Deaths asso
ciated with anaesthesia Cases in which hypotension has been used
Anaesthesia 8 263 1953
- 85 COSTA J and TORRES R P Comentarios a cien adenomectomias
prostáticas Empleo de la hipotension controlada *Arch españ
urol* 8 331 1952
- 86 ————— Hipotensao controlada e protatectomia retro
pública (nota prévia) *Portugal méd* 36 125 1952
- 87 ————— Vantagens da hipotensão controlada na
adenomectomia protática *Portugal méd* 36 585 1952
- 88 COTTEN M DE V BROWN J M and KRONEN P S Heart force
responses to pressor amines during hypotension produced by hex
amethonium *Anesthesiology* 15 126 1954
- 89 CO TUI BURSTEIN C L and RUGGIERO W F Total spinal block a
preliminary report *Anesthesiology* 1 280 1940
- 90 CRILE G W and LOWER W E *Anoci association* Philadelphia
W B Saunders Co 1914

- 91 CROSETTI J P MULLER C A and PETTAVEL J Le traitement des embolies pulmonaires par les ganglioplegiques *Presse méd*, 60 1529 1952
- 92 CRUMPTON C W and MURPHY Q R Effects of hexamethonium bromide upon hemodynamics of cerebral and coronary circulation in hypertension *J Clin Investigation* 31 622 1952
- 93 CUSACK P B and CUSACK A L J Hexamethonium bromide in vascular hypertension *J Irish M.A* 28 47 1951
- 94 DALE H H and LAIDLAW P E Histaminic shock *J Physiol* 52 355 1919
- 95 DANZIG L E and GOMEZ A C Long term intravenous use of arfonad in the treatment of pulmonary edema *Am J M Sc* 228 626 1954
- 96 DAUMET F SAUVAGE Y and ANGLES C A propos de cent résections pulmonaires réalisées sous hypotension contrôlée *Poumon*, 8 777 1952
- 97 DAVIES R M MOORE F T and WYNN WILLIAMS D Treatment of acute pancreatitis with hexamethonium bromide *Brit.M J* 2 1251 1953
- 98 DAVISON M H A Pentamethonium iodide in anaesthesia *Lancet* 1 252 1950
- 99 ——— Hypotension *Anaesthesia* 7 116 1952
- 100 ——— Death after anaesthetic with hypotension *Lancet* 2 1273 1952
- 101 ——— The case against hypotension *Anaesthesia* 8 255 1953
- 102 DE CLIVE LOWE S G and NORTH J Controlled hypotension with arfonad *Lancet* 1 680 1954
- 103 DEWAR H A OWEN S G and JENKINS A R Effect of hexamethonium bromide on the cerebral circulation in hypertension *Brit.M J* 2 1017 1953
- 104 DE WARDENER H E MILES B E WYLIE W D and CHURCHILL-DAVIDSON H C Complications of controlled hypotension *Lancet* 2 93 1952
- 105 DAURI A L uso degli ipotensivi in chirurgia con particolare riguardo alla chirurgia toracica *Minerva anest* 19 113 1953
- 106 DOCAL J C Hipotension controlada con novocaina y pendiomid *Dis med B Air* 26 21 1954
- 107 DONATI G S and PERUZZO L Interventi di chirurgia generale in ipotensione controllata *Acta anaesth Padova* 3 217 1952
- 108 DOUTHWAITE, A H and THORNE M G The effects of hexamethonium bromide on the stomach *Brit.M J* 1 111 1951
- 109 DUBOUCHER H and SEROR J Traitement des hémorragies graves par l'hypotension contrôlée *Mém Acad chir* 78 934 1952

- 110 DU BOUCHET N LATSCHA B I and PASSALECQ J Premiers résultats de l'usage de l'hexaméthonium au cours des interventions de chirurgie du cœur *Anesth et analg* 9 361 1952
- 111 DU CAILLAR J La prévention du choc opératoire par les méthoniums en chirurgie viscérale a propos de 120 observations *Presse méd* 61 1595 1953
- 112 ——— and ROGER M La place des méthoniums dans les associations végétativolytiques a propos de 140 observations *Anesth et analg* 10 435 1953
- 113 DUNDEE J W FRANCIS I L and SEDZIMIR C B Safer hypotension *Lancet* 1 885 1954
- 114 EBDLI C Lipotensione controllata da anestesia peridurale *Minerva anest* 19 125 1953
- 115 ECKENHOFF J E HAFKENSCHIEL J H FOLTZ E L and DRIVER, R L Influence of hypotension on coronary blood flow cardiac work and cardiac efficiency *Am J Physiol* 152 545 1948
- 116 EDDLEMAN E E JR WILLIS K and HEYER H E The effect of posture on the cardiac cycle posteroanterior cardiac diameters and apparent stroke volume as studied by the electrokymograph *Am Heart J* 40 504 1950
- 117 EDITORIAL Methonium controlled hypotension in surgery *J.A.M.A* 151 129 1953
- 118 ——— Controlled hypotension *Post Grad MJ* 28 331 1952
- 119 EMMETT J L Subarachnoid injections of procaine hydrochloride quantitative effects of clinical doses on sensory sympathetic and motor nerves *J.A.M.A* 102 425 1934
- 120 ENDERBY G E H Controlled circulation with hypotensive drugs and posture to reduce bleeding in surgery Preliminary results with pentamethonium iodide *Lancet* 1 1145 1950
- 121 ——— Controlled hypotension *Lancet* 2 145 1952
- 122 ——— Hypotensive anaesthesia in surgery *Ann Roy Coll Surgeon England* 11 310 1952
- 123 ——— Hypotensive drugs in anaesthesia *Brit J Plast Surg*, 5 221 1953
- 124 ——— Postural ischaemia and the use of the methonium compounds *Anesth & Analg* 32 1 1953
- 125 ——— Postural ischaemia and blood pressure *Lancet* 1 185 1954
- 126 ——— and PELMORE J F Controlled hypotension and postural ischaemia to reduce bleeding in surgery Review of 250 cases *Lancet* 1 663 1951
- 127 ——— DAVISON M H A KORKIS F B BARNES M SHACKLETON R P W SCURR C F GILLIES J MUSGROVE A. H and WYMAN J B Discussion on the use of hypotensive drugs in surgery *Proc Roy Soc Med* 44 829 1951

- 128 ETSTEN H YORK G E and HAWWICH H E Pattern of metabolic depression induced with pentothal sodium *AMA Arch Neurol & Psychiat* 56 171 1946
- 129 EVANS H and ENDERBY G E II Controlled hypotension and its effect on renal function *Lancet* 1 1045 1952
- 130 FANJEUX O and SOULIER J A propos de quelques observations d'anesthésies sous hibernation et sous hypotension contrôlée *Afrique fr chir* 3-4 112 1952
- 131 FERGUSON L K and NORTH J P Observations on experimental spinal anesthesia *Surg Gynec & Obst*, 54 621 1932
- 132 FERUGLIO F S ROMERO C and RUIV P II flusso sanguigno cerebrale ed il consumo di O_2 nell'ipotensione da ganglioplegici *Mis nerva med* 45 1422 1954
- 133 FIBAK J Magen speiserohren blutung aufgehalten durch kontrol lierte blutdrucksenkung mit reichlicher bluttransfusion *Schweiz med Wchnschr* 84 39 1954
- 134 ——— and KIELBOWICZ D Sztuczne obnizanie cisnienia krwi celem zmniejszenia krwawienia podczas zabiegow chirurgicznych *Polski tygodnik lek* 9 65 1954
- 135 FINE J BENTEL H and MEYEROVITZ L Controlled hypotension in fenestration operations experiences in twenty seven cases *South African MJ* 28 101 1954
- 136 FINNERTY F A JR and FREIS E D Experimental and clinical evaluation in man of hexamethonium (C6) a new ganglionic blocking agent *Circulation* 2 828 1950
- 137 ——— Clinical appraisal of hexamethonium (C6) in peripheral vascular diseases *New England J Med* 245 325 1951
- 138 FINNERTY F A JR WITKIN L and FAZEKAS J F Acute hypotension and cerebral hemodynamics preliminary report *M Ann District of Columbia* 22 115 1953
- 139 FORD R V MOYER J H and SPURR C L Hexamethonium in the chronic treatment of hypertension—its effect on renal hemodynamics and on the excretion of water and electrolytes *J Clin Investigation* 32 1133 1953
- 140 FRANKEL E Methonium compounds in hypertension *Lancet* 1 408 1951
- 141 FREEMAN N E SHAFFER S A SCHECTER A E and HOLLING H E The effect of total sympathectomy on the occurrence of shock from hemorrhage *J Clin Investigation* 17 359 1938
- 142 FREIS E D Methonium compounds in hypertension *Lancet* 1 909 1951
- 143 ——— Veratrum viride and hexamethonium in the treatment of severe diastolic hypertension *M Ann District of Columbia* 20-297 1951

- 144 ——— FINNERTY F A JR SCHNAPER H W and JOHNSON R L The treatment of hypertension with hexamethonium *Circulation* 5:20 1952
- 145 ——— ROSE J C HICCINS T F KELLEY R T SCHNAPER H W and JOHNSON R L The hemodynamic effects of hexamethonium in man *J Clin Investigation* 31:629 1952
- 146 ——— SCHNAPER H W JOHNSON R L ROSE J C and FINNERTY F A JR Clinical evaluation of hexamethonium a new ganglionic blocking agent—preliminary report *Am J Med* 11:242 1951
- 147 ——— STANTON J R FINNERTY F A JR SCHNAPER H W JOHNSON R L ROTH C E and WILKINS R W The collapse produced by venous congestion of the extremities or by venesection following certain hypotensive agents *J Clin Investigation* 30:435 1951
- 148 FREY R Fortschritte und erfahrungen mit der künstlichen blut drucksenkung *Arch klin Chir* 276:670 1953
- 149 GARDNER W J The control of bleeding during operation by induced hypotension *JAMA* 132:572 1946
- 150 ——— and HALE D E Arterial bloodletting during operation as aid in hemostasis *Am J Surg* 79:635 1950
- 151 ——— and LING A Controlled hypotension by the bleeding method in operations for intracranial meningiomas *Surg Gynec & Obst* 98:343 1954
- 152 GASSER H S and ERLANGER J Role of fiber size in the establishment of a nerve block by pressure or cocaine *Am J Physiol* 88:581 1929
- 153 GENAZZANI E I ganglioplegia in medicina e chirurgia con particolare riguardo alla ipotensione controllata *Riforma med* 67:914 1953
- 154 ——— and RUSSO A Contributo sperimentale all azione dei ganglioplegia impiegati per ipotensione controllata *Gior ital chir* 9:547 1953
- 155 GERLACH H Gesteuerte blutdrucksenkung durch ganglionblocker (pendiomid) in der operativen otorhino laryngologie *Arch Ohren Nasen u Kehlkopf* 162:463 1953
- 156 GERTNER S B LITTLE D M JR and BONNYCASTLE D D The urinary excretion of arfonad by patients undergoing controlled hypotension during surgery *Anesthesiology* (in Press)
- 157 GILI G Lipotensione controllata in chirurgia *Minerva chir.* 8:288 1953
- 158 GILLAN J G Two cases of unilateral blindness following anaesthesia with vascular hypotension *Canad MAJ* 69:294 1953
- 159 GILLIES J Anaesthesia for the surgical treatment of hypertension *Proc Roy Soc Med* 42:295 1949

- 160 ——— Anaesthetic factors in the causation and prevention of excessive bleeding during surgical operations *Ann Roy Coll Surgeons England* 7:204 1950
- 161 ——— Physiological trespass in anaesthesia *Proc Roy Soc Med* 45:1 1952
- 162 GILMORE H R KOPELMAN H McMICHAEL J and MILNE, I G The effect of hexamethonium bromide on the cardiac output and pulmonary circulation *Lancet* 2:898 1952
- 163 GLASSER O and PAGE I H Prognostic signs in experimental hemorrhagic shock *Cleveland Clin Quart*, 13:125 1946
- 164 ——— Intra arterial transfusion in the treatment of experimental hemorrhagic shock *Cleveland Clin Quart* 14:121 1947
- 165 ——— Experimental hemorrhagic shock a study of its production and treatment *Am J Physiol* 154:297 1948
- 166 GLENN W W L HAMPTON L J and GOODYER A V N Use of controlled hypotension in large blood vessel surgery *A.M.A Arch Surg* 68:1 1954
- 167 GNUCHTEL W Ergebnisse der fortlaufenden temperaturmessung bei anaesthesie künstlicher blutdrucksenkung und hypothermie *Anaesthesist* 3:80 1954
- 168 GOFFEN H S FITZPATRICK L J and WOOD P M Control of bleeding during fenestration and similar operations by sympathetic block anesthesia *Anesth & Analg* 29:340 1950
- 169 GOLDSMITH A J H and HEWER A J H Unilateral amaurosis with partial recovery after using hexamethonium iodide *Brit.M.J.* 2:759 1952
- 170 GOLDSTONE H Death associated with methonium treatment *South African M.J.* 26:552 1952
- 171 GRABNER G MLCZOK F STEINBEREITHNER, K and VETTER H Der kreislauf bei künstlicher blutdrucksenkung *Anaesthesist* 3:50 1954
- 172 GRAHAM A J P and DOUGLAS D M Effect of the head down position on the circulation in hypotensive states *Lancet* 2:941 1949
- 173 GREENE N M Hypotensive spinal anesthesia *Surg Gynec & Obst*, 95:331 1952
- 174 ——— BUNKER J P KERR W S VON FELSINGER J M KELLER J W and BEECHER H K Hypotensive spinal anesthesia respiratory metabolic hepatic renal and cerebral effects *Ann Surg* 140:641 1954
- 175 GRIFFITHS H W C and GILLIES J Thoraco lumbar splanchnicectomy and sympathectomy anaesthetic procedure *Anaesthesia* 3:134 1948
- 176 GRIFFITHS J A The effects of general anaesthesia and hexamethonium on the blood sugar in non diabetic and diabetic surgical patients *Quart J Med* 22:405 1953

- 177 GROB H and HARVEY A M Observations on the effects of the autonomic blocking agent bis trimethylammonium pentane dibromide (C5) in normal subjects and in patients with peripheral vascular disease and hypertension and comparison with tetraethylammonium chloride *Bull Johns Hopkins Hosp* 87 616 1950
- 178 GROB D and LANGFORD H G Further observations on the effects of autonomic blocking agents in patients with hypertension I General systemic effects of hexamethonium pentamethonium and hydrazinophthalazine *Circulation* 8 205 1953
- 179 GROB D SCARBOROUGH W R KATTUS A A JR and LANGFORD H G Further observations on the effects of autonomic blocking agents in patients with hypertension II Hemodynamic ballistocardiographic and electrocardiographic effects of hexamethonium and pentamethonium *Circulation* 8 352 1953
- 180 GROVES L A EFFLER D B and SOVES F M JR Controlled hypotension in the surgical treatment of certain cases of patent ductus arteriosus *Cleveland Clin Quart* 21 169 1954
- 181 GUGLIELMI F Gli steroidi urinari negli operati con narcosi in ipotensione *Gior ital anestesiol* 18 512 1952
- 182 ——— and BELSASSO E Le variazioni del metabolismo dopo ganglioplegici *Gior ital anestesiol* 18 530 1952
- 183 ——— and BIACINI S Le variazioni glicemiche nell'anestesia con ipotensione *Gior ital anestesiol* 18 523 1952
- 184 ——— and PIZZIOLI R Modificazioni ematiche nell'anestesia con ipotensione posturale *Gior ital anestesiol* 18 499 1952
- 185 ——— RAMOVECCHI E and RICCI C La velocità di circolo e la pressione venosa sotto l'azione dei ganglioplegici *Gior ital anestesiol* 18 517 1952
- 186 GUIDETTI B and ALVISEI C Lesametonio in neuro chirurgia e in neurologia *Riv pat nerv* 73 119 1954
- 187 GUIOT G L'hypotension contrôlée par les ganglioplégiques ses indications en neuro chirurgie *Gaz med France* 59 1441 1952
- 188 ——— and DAMOISEAU B Vers la suppression des hémorragies et de l'œdème cérébral par l'emploi des ganglioplégiques en neuro chirurgie 24 observations *Anesth et analg* 8 641 1951
- 189 ——— Des possibilités d'une neuro chirurgie exsangue et sans œdème cérébral grâce à l'emploi des substances ganglioplegiques *Rev prat Paris* 2 172 1952
- 190 GURD F B VINEBERG A M and BOURNE W Our experiences in the employment of spinal anesthesia for thoracoplasty *J Thoracic Surg* 7 506 1938
- 191 ——— Further experiences in the use of spinal anesthesia for thoracoplasty *Ann Surg* 110 872 1939
- 192 HABIB D V PAPPER E M FITZPATRICK H F LOWRANCE P

- SMYTHIE C M and BRADLEY S E Renal and hepatic blood flow glomerular filtration rate and urinary output of electrolytes during cyclopropane ether and thiopental anesthesia operation and immediate post operative period *Surgery* 30 241 1951
- 193 HAD II Anaesthesien in gesteuerter blutdrucksenkung *Arch klin Chir* 276 686 1953
- 194 ——— Erfahrungen mit der künstlichen blutdrucksenkung *Anaesthesist* 2 III 1953
- 195 ——— Klinik und "low pressure" mit besonderer berücksichtigung von EKG veränderungen *Anaesthesist* 3 49 1954
- 196 HALE D E Controlled hypotension by arterial bleeding during operation and anesthesia *Anesthesiology* 9 498 1948
- 197 HAMPTON L J and LITTLE D M JR Results of a questionnaire concerning controlled hypotension in anaesthesia *Lancet* 1 1299 1953
- 198 ——— Complications associated with the use of controlled hypotension in anesthesia *A.M.A Arch Surg* 67 549 1953
- 199 HARPER J A Pentamethonium in surgical emergencies *Brit.M.J* 1 1262 1951
- 200 HARRIS H E and HALE D E Induced hypotension in control of bleeding during fenestration operation *Cleveland Clin Quart* 14 159 1947
- 201 HAY E B and WEBB J A The effect of increased arterial blood flow to the liver on the mortality rate following hemorrhagic shock *Surgery* 29 826 1951
- 202 HAYNES B W JR MORRIS G C JR MOYER J H and SNYDER H B The effects of controlled hypotension on glomerular filtration and renal plasma flow in man *Surg Forum* 4 443 1953
- 203 HAYWARD G W Anaesthesia and the cardiac patient *Anaesthesia* 7 67 1952
- 204 HEINBECKER P BISHOP G H and O'LEARY J L Analysis of sensation in terms of the nerve impulse *A.M.A Arch Neurol & Psychiat* 31 34 1934
- 205 HERMANN H MORIN G and VIAL J Les régulations périphériques chez le chien sous moelle évolution de la pression artérielle au cours de la destruction ascendante progressive de la moelle épinière chez le chien *Arch internat de physiol* 43 23 1936
- 206 HERSHEY S G ZWEIFACH B W and METZ D B An evaluation of the protective action of autonomic blocking agents in peripheral circulatory stress *Anesthesiology* 15 589 1954
- 207 HEWER A J H Some observations on induced hypotension with special reference to its use in neurosurgery *Arch Middlesex Hosp* 3 189 1953

- 208 HEYMANS C *Introduction to the Regulation of the Blood Pressure and Heart Rate* Springfield Illinois Charles C Thomas Publisher 1950
- 209 HICKAM J B and PRYOR W W Cardiac output in postural hypotension *J Clin Investigation* 30 401 1951
- 210 HIZENERT G and KUCHER R Transventrikuläre pulmonalstenosen operation in ganglienblockade mit kontrollierter blutdrucksenkung *Wien klin Wchnschr* 63 198 1953
- 211 HILL K R and GOODWIN M P Influence of hypotension on the intracapsular cataract operation *Brit J Ophth* 36 683 1952
- 212 HILLIS H R and KELLY J C C Hiccup and hexamethonium *Brit M J* 1 1143 1951
- 213 HIRSON C and KELSALL A R Methonium compounds in hypertension *Lancet* 1 585 1951
- 214 HOFFMANN T Die herabsetzung des peroperativen blutverlustes durch hexamethonium und neigungslagerung *München med Wchnschr* 94 711 1952
- 215 ——— Die anwendung von blutdrucksenkenden mitteln in der lungenchirurgie (künstliche blutdrucksenkung) *Arch klin Chir* 276 690 1953
- 216 HOLLE F and KERN E Über das verhalten der nierenfunktion nach gesteuerter blutdrucksenkung *Klin Wchnschr* 32 67 1954
- 217 HOLT M C and LITCHFIELD J W Bromism on low salt diets *Lancet* 1 347 1951
- 218 HOPKIN D A B Safer hypotension *Lancet* 1 1186 1954
- 219 HORATZ K Normale und paradoxe kreislaufreaktionen bei der künstlichen blutdrucksenkung *Anaesthetist* 2 169 1953
- 220 ——— and STURTZBECHER F Neue hilfsmittel in der anästhesie *Anaesthetist* 1 149 1952
- 221 HUEBER E F SAEKINGER H and WOHLRAUB K Zur kreislauf mechanismus bei der künstlichen blutdrucksenkung *Anaesthetist* 3 55 1954
- 222 HUET P C LABAYLE J and ROTH F Fibrome nasopharyngien opere sous hypotension contrôlée *Ann oto laryng* 70 792 1953
- 223 ——— A propos de l'emploi des méthoniums en cancérologie cervico faciale *Semaine hôp Paris* 29 2868 1953
- 224 HUGHES G Use of pentamethonium and posture to reduce bleeding in fenestration operations *Lancet* 1 666 1951
- 225 ——— Controlled hypotension *Lancet* 2 145 1952
- 226 ——— Postural ischaemia and blood pressure *Lancet* 1 311 1954
- 227 HUNTER A R Hexamethonium bromide *Lancet* 1 251 1950
- 228 ——— and HEWER A J H The present position of anaesthesia for neurosurgery *Proc Roy Soc Med* 45 427 1952

- 229 JACKSON I Arteriotomy in neurosurgery *Anaesthesia* 9 13 1954
- 230 JAMES A COULTER R L and SAUNDERS J W Controlled hypotension in neurosurgery With special reference to hypotension induced by pneumatic suction applied to the legs *Lancet* 1 412 1953
- 231 JENKIN P S Control of surgical bleeding by the production of hypotensive states preliminary report *South African M J* 25 842 1951
- 232 JONES R R Hypotensive anesthesia *Med Bull U S Army Europe* 11 127 1954
- 233 JONNESCO T Remarks on general spinal analgesia *Brit.M J* 2 1396 1909
- 234 JUNGLENG O Kontrollierte blutdrucksenkung mit TEAB *Zentralbl Chir.* 78 1887 1953
- 235 JUST O Die vermindernng des operativen blutverlustes durch kunstliche hypotension *Arztl Wchnschr* 7 433 1952
- 236 KAPLAN S A and ASSALI N S Effects of apresoline veratrum alkaloids high spinal anesthesia and arfonad on renal hemodynamics of pregnant patients with toxemia and essential hypertension *Surg Gynec & Obst* 97 501 1953
- 237 KAY A W and SMITH A N Effect of hexamethonium iodide on gastric secretion and motility *Brit.M J* 1 460 1950
- 238 ————— Effect of oral hexamethonium salts on gastric secretion *Brit.M J* 2 807 1950
- 239 KENNY S Controlled hypotension *Irish J M Sc* 6 241 1952
- 240 KERN E Lanesthésie pour la sympathectomie dorsale haute et la sympathectomie dorso lombaire (opérations de Smithwick) *Anesth et analg* 7 481 1950
- 241 ————— Blutungsverminderung durch gesteuerte herabsetzung des blutdruckes *Anaesthesist* 1 15 1952
- 242 ————— Problems of anaesthesia and transfusion in major surgery on the hip joint *Anesth & Analg* 31 350 1952
- 243 ————— L'hypotension contrôlée une technique pour diminuer le saignement chirurgical *Semaine méd Paris* 28 646 1952
- 244 ————— La place de l'hypotension contrôlée dans la pratique courante *Anesth et analg* 10 315 1953
- 245 ————— Die praxis der kunstlichen blutdrucksenkung nach 2 jahren klinischer erfahrung *Arch klin Chir* 276 685 1953
- 246 ————— and MENNUNI R La diminuzione dell emorragia medi ante la tecnica dell ipotensione controllata osservazioni su 130 casi *Gior ital chir* 8 401 1952
- 247 ————— CHATEAUREYNAUD J LASSNER J and VOURCH G La nostra esperienza dell ipotensione controllata osservazioni su 416 casi *Gior ital anestesiol* 18 545 1952
- 248 ————— SCHWARTZ M BATAILLE L and ESPINASSE J Analyse

- de 166 observations d'hypotension contrôlée *Anesth et analg* 9 327 1952
- 249 KETY S S and SCHMIDT C F The determination of cerebral blood flow in man by the use of nitrous oxide in low concentrations *Am J Physiol* 143 53 1945
- 250 ————— Nitrous oxide method for quantitative determination of cerebral blood flow in man theory procedure and normal values *J Clin Investigation* 27 476 1948
- 251 KILBUFF C J The use of arfonad in controlled hypotension *Lancet* 1 337 1954
- 252 KILPATRICK J A and SMIRK F H Comparison of oral and subcutaneous administration of methonium salts in the treatment of high blood pressure *Lancet* 1 8 1952
- 253 KIRKENDALL W M and CULBERTSON J W Some effects of hexamethonium on renal circulation *J Clin Investigation* 31 644 1952
- 254 KIRSCH J Blutstillung durch ganglienblockade (pendiomid) *Med Klin Berlin* 48 1516 1953
- 255 KOHLSTAEDT K G and PAGE I H Terminal hemorrhagic shock. Circulatory dynamics recognition and treatment *Surgery* 16 450 1944
- 256 KOLBL H KUCHER R and STEINBEREITHNER K Die veränderungen der bluteiweisskörper unter künstlicher blutdrucksenkung *Anaesthetist* 3 56 1954
- 257 KORKIS F B Controlled hypotension in surgery *Lancet* 1 801 1951
- 258 KOSTER H Spinal anesthesia with special reference to its use in surgery of the head neck and thorax *Am J Surg* 5 554 1928
- 259 ————— and KASMAN L E Spinal anesthesia for the head neck and thorax its relation to respiratory paralysis *Surg Gynec & Obst* 49 617 1929
- 260 KUCHER R Die operation unter kontrollierter blutdrucksenkung *Schweiz med Wchnschr* 82 1219 1952
- 261 ————— and STEINBEREITHNER K Funktionell pathologische gesichtspunkte zum indikationsproblem der beabsichtigten blutdrucksenkung *Anaesthetist* 2 167 1953
- 262 ————— Über einen neuen ganglienblocker *Anaesthetist* 2 168 1953
- 263 ————— Bemerkungen zu umstrittenen problemen der künstlichen blutdrucksenkung *Anaesthetist* 3 47 1954
- 264 KUHLMANN J and TRAUTMANN R L'hypotension contrôlée en chirurgie thyroïdienne *Strasbourg méd* 5 302 1954
- 265 LABORIT H Sur le mécanisme d'action des méthoniums *Anesth et analg* 9 386 1952
- 266 ————— Sul meccanismo dell'ipotesione controllata fisiologia e mezzi farmacologici per ottenerla *Gior ital anestesiol* 18 541 1952

- 267 LANDIS E M Capillary blood pressure in mammalian mesentery ■
determined by micro injection method *Am J Physiol* 93 353
1930
- 268 LANG H Kontrollierte blutdrucksenkung und periduralanästhesie
Chirurgia 25 181 1954
- 269 LANGSTON A W Controlled hypotension to reduce surgical haemor-
rhage *Canad MAJ* 69 371 1953
- 270 LASSNER J La diminution du saignement par la technique de
l'hypotension contrôlée (Étude de 72 observations dont 45 en
chirurgie pelvienne) *Anesth et analg* 9 341 1952
- 271 ——— Contribution à l'étude des effets du pentaméthonium
Anesth et analg 9 347 1952
- 272 LAURENCE B R and STACEY R S Hypoglycaemia masked by
methonium *Lancet* 2 1145 1951
- 273 ——— Effect of hexamethonium on the response to
insulin in animals and man *Brit J Pharmacol* 7 255 1952
- 274 LAZORTHES G CAMPAN L ANDUZE H and STELLET L L'hypoten-
sion contrôlée par les dérivés du méthonium pratique chirurgicale
Presse méd 60 1253 1952
- 275 LEADING ARTICLE Complications of controlled hypotension *Lancet*
2 25 1952
- 276 ——— Deliberate hypotension during anaesthesia *M J Austra-
lia* 2 517 1952
- 277 LEARMONTH J The surgeon and ischaemia *Brit.MJ* 1 743 1953
- 278 LEATHART G L Postural ischaemia and blood pressure *Lancet*
1 311 1954
- 279 LE BRIGAND J Suites opératoires observées chez 24 malades soumis
à l'hexaméthonium *Anesth et analg* 9 392 1952
- 280 LEDRU R Essai de traitement d'une hémoptysie anormalement pro-
longée et rebelle par l'hypotension contrôlée prolongée *Lyon
méd* 189 12 1953
- 281 LE FILLIATRE G *Précis de rachianesthésie générale* Paris Libr Le
Francois 1921
- 282 LEMOYNE J Fibrome naso pharyngien et hypotension contrôlée
Ann oto laryng 70 565 1953
- 283 LEONHARDT K O Übersichten Blutsparendes operieren durch
künstliche blutdrucksenkung *Anaesthesist* 1 116 1952
- 284 LEVEEN H H MULDER A G and PROKOP F The physiological
mechanism for death in massively bleeding peptic ulcer *Surg
Gynec & Obst* 94 433 1952
- 285 LEWIS I Value of hypotensive drugs in minimizing blood loss in
thoracic surgery *Lancet* 2 150 1951
- 286 LITTLE H M JR HAMPTON L J and GROSSKREUTZ D C The use
of arfonad in a modification of the technique of controlled hypo-
tension *Surgery* 35 857 1954

- 287 LOCKET S Methonium compounds in hypertension *Lancet*, 1 473 1951
- 288 ——— Controlled hypotension and its effect on renal function *Lancet* 1 1166 1952
- 289 ——— SWANN P G and GRIEVE W S M Methonium compounds in the treatment of hypertension *Brit.M.J.* 1 778 1951
- 290 LOEW F Erfahrungen mit der kontrollierten blutdrucksenkung bei intrakraniellen operationen *Arch klin Chir* 276 694 1953
- 291 LONGTIN L FOURNIER J MIGNAULT G LONGTIN J and JOLY R Rapport préliminaire de l'étude de 25 cas d'hypotension contrôlée en chirurgie thoracique *Union méd Canada* 81 390 1952
- 292 LONGTIN J LONGTIN L BOUSQUET E and MÉTHOT J P Hypotension contrôlée en chirurgie thoracique étude clinique et électrocardiographique de 100 cas et de 100 témoins *Union méd Canada* 83 625 1954
- 293 LORIAN P H and MILLER M I Clinical experience with hypotensive anesthesia *Am Surgeon* 20 629 1954
- 294 LOTTENBACH K Die wirkung des N N N N 3 pentamethyl N,N diethyl 3 aza pentan 1 5 diammonium dibromid (Ciba 9290) auf die kreislaufreflexe beim carotissinussyndrom *Schweiz med Wchnschr* 81 310 1951
- 295 LOVE J G COURTIN R F ANDERSON C D and LUNDY J S Hypotension induced by arteriotomy using Pages technic an aid in surgery its use in the removal of an epidermoid tumor of the third ventricle report of case *Proc Staff Meet Mayo Clinic* 23 153 1948
- 296 LYNN R B SANCETTA M and SIMEONE F A Studies of hemodynamic changes in human beings following induction of low and high spinal anesthesia *J Lab & Clin Med* 38 924 1951
- 297 ——— ——— ——— and SCOTT R W Observations on the circulation in high spinal anesthesia *Surgery* 32 195 1952
- 298 LYNN R B and SHACKMAN R The peripheral circulation during general anaesthesia and surgery *Brit.M.J.* 2 333 1951
- 299 McCORMICK V The use of hypotension in surgery *J Roy Nav M Serv* 40 6 1954
- 300 McCUBBIN J W and PAGE I H Nature of the hypotensive action of a thiophanium derivative (RO 2 2222) in dogs *J Pharmacol & Exper Therap* 105 437 1952
- 301 MCGILL R J Hiccup and hexamethonium *Brit.M.J.* 1 948 1951
- 302 ——— Methonium in intractable hiccup *Lancet* 1 1018 1951
- 303 McLAUGHLIN C R and WATSON J Operative aspects of postural ischaemia with hexamethonium bromide *Brit J Plast Surg* 4 121 1951

- 304 McMICHAEL, J The oxygen supply of the liver *Quart J Exper Physiol* 27 73 1937
- 305 McNALLY P A Some results of treatment with hexamethonium bromide in cases of vascular hypertension *J Irish M A.* 28 44 1951
- 306 McQUEEN E. G. Hexamethonium bromide and kidney function *M J Australia* 1769 1952
- 307 MACKEY W A Hexamethonium in hypertension *Brit.M J* 1 918 1951
- 308 MACKINNON J Effect of hypotension producing drugs on the renal circulation *Lancet* 2 12 1952
- 309 MACILL I W Recent developments in anaesthesia in the British Isles with particular reference to induced hypotension *Anesthesiology* 15 103 1954
- 310 ——— Scurr C. F and WYMAN J B Controlled hypotension by a thiophanium derivative. *Lancet* 1 219 1953
- 311 MANDOW G A ARDEN G P and STONEHAM F J R Hypotension induced with hexamethonium bromide during anaesthesia *Brit J Anaesth* 26 26 1954
- 312 MANNI C Lipotensione controllata quale mezzo di riduzione delle emorragie intraoperatorie *Policlinico se. prat* 59 765 1952
- 313 MARLAND T J The present status of the hypotensive drugs in surgery *Mississippi Doctor* 30 240 1953
- 314 MARKER, A and MIESCHER K. Über die stufenweise quaternisierung von aliphatischen polyaminen Neue verbindungen mit ganglien blockierender wirkung *Helvet chim acta* 34 924 1951
- 315 MASON A A and PELMORE J F Combined use of hexamethonium bromide and procaine amide in controlled hypotension a preliminary report. *Brit.M J* 1 250 1953
- 316 MATHEY J and BIVET J F L'hypotension contrôlée et l'usage des sels du méthonium en chirurgie *Rev prat* 2 1731 1952
- 317 MENDELSON, M L and SATU C Relationship of renal function to blood pressure during ganglionic blockade in the anesthetized dog *Am J Physiol.* 173 355 1953
- 318 MENEZES L. C. Anestesia com hipotensão induzida em neurocirurgia *Med cir farm Rio* 203 362 1953
- 319 MERLE H AUBICNE R and KERN E Hypotension contrôlée rapport préliminaire *Anesth et analg* 8 635 1951
- 320 ——— Bilan de deux années d'expérience de l'hypotension contrôlée *Mém Acad chir* 79 368 1953
- 321 MILES H E DE WARDENER H E. CHURCHILL DAVIDSON H C. and WYLIE W D. Effect on renal circulation of pentamethonium bromide during anaesthesia. *Clin Sc* 11 73 1952

- 322 MILETTI M and SURIANI U Controlled hypotension with ganglionic blocking agents in neurosurgery *Acta neurochir* 3:275 1954
- 323 MILLS L C and MOYER J H The acute effects of hexamethonium on renal hemodynamics in normotensive and hypertensive human subjects *Am J M Sc* 226:1, 1953
- 324 MILNE G E and OLFESKY S Excretion of the methonium compounds *Lancet* 1:889 1951
- 325 VOLFETTA N and MANGIAVACCHI E Ipotensione controllata e prostatectomia retropubica *Minerva anest* 19:72 1953
- 326 MONDRVI P and FABRIS F Ganglioplegia e shock emorragico ricerche sperimentali e deduzioni cliniche *Acta anaesth Padova* 4:157 1953
- 327 MONTAG H Blutsparendes operieren in gesteuerter blutdrucksenkung *Zentralbl Chir* 78:623 1954
- 328 MOON V H *Shock Its Dynamics Occurrence and Management* Philadelphia Lea & Febiger 1952
- 329 MORICCA G Ipotensione controllata con arfonad (RO 2 2222) con particolare riguardo agli interventi di neurochirurgia ricerche cliniche e sperimentali *Rassegna Internaz clin e terap* 31:362 1954
- 330 MORRIS N Hexamethonium compounds in the treatment of pre-eclampsia and essential hypertension during pregnancy *Lancet* 1:322 1953
- 331 MORRIS G C JR MOYER J H SNYDER H B and HAYNES M W JR Cerebral hemodynamics in controlled hypotension *Surg Forum* 4:140 1953
- 332 _____ Vascular dynamics in controlled hypotension a study of cerebral and renal hemodynamics and blood volume changes *Ann Surg* 138:706 1953
- 333 MORTIMER P L F A new apparatus for the production of controlled hypotension by arteriotomy *Anaesthesia* 6:128 1951
- 334 MORTON A W Subarachnoid injection of cocaine *Pacific M J* 43:802 1950
- 335 MOYER J H HUGGINS P A HANDLEY C A and MILLS L C Effect of hexamethonium chloride on cardiovascular and renal hemodynamics and on electrolyte excretion *J Pharmacol & Exper Therap*, 106:157 1952
- 336 MOYER J H and MILLS L C Hexamethonium—its effect on glomerular filtration rate maximal tubular function and renal excretion of electrolytes *J Clin Investigation* 32:172 1953
- 337 MOYER J H and MORRIS G C JR Cerebral hemodynamics during controlled hypotension induced by the continuous infusion of ganglionic blocking agents (hexamethonium pendiomide and arfonad) *J Clin Investigation* 33:1081 1954

- 338 MUELLER R P LYNN R B and SANCETTA S M Studies of hemodynamic changes in humans following induction of low and high spinal anesthesia II The changes in splanchnic blood flow oxygen extraction and consumption and splanchnic vascular resistance in humans not undergoing surgery *Circulation* 6 891 1952
- 339 MUKHERJEE S H and ROWLANDS S Pulmonary and peripheral circulation times in haemorrhagic shock *Lancet* 1 1041 1951
- 340 MUNDLEER, P PROOT R and REMOUCHANGES L Essais de saignement contrôlée par un hypotenseur *Anesth et analg*, 9 322 1952
- 341 MURPHY E. A Treatment of hypertension with hexamethonium bromide *Lancet* 2 899 1951
- 342 NEEDLEMAN H L. and HORWITZ O. A comparative study of the effects of three vasodilator drugs pentamethonium bromide (C5) dilatol (SKF 1700 A) and pendiomide (BA 929c) on the digital cutaneous flow *Am J M Sc.* 226 161 1953
- 343 NEUMANN C FOSTER, A D JR. and ROVENSTINE E. A. The importance of compensatory vasoconstriction in unanesthetized areas in the maintenance of blood pressure during spinal anesthesia *J Clin Investigation* 24 345 1945
- 344 NICHOLSON M J SARNOFF S J and CREHAN J P The intravenous use of a thiophanium derivative (arfonad—RO 2 222) for the production of flexible and rapidly reversible hypotension during surgery *Anesthesiology* 14 215 1953
- 345 NILSSON F The application of a method for the investigation of cerebral damage following anaesthesia using controlled hypotension a preliminary communication *Brit J Anaesth* 25 24 1953
- 346 NORPOTH L DRUGEMOLLER P and FLOSSDORF T Behandlung des herpes zoster mit dem ganglien blockierenden präparat pendiomid (9295 Ciba) *Deutsche med Wchnschr* 76 1066 1951
- 347 O'BRIEN D and COLE W H J The use of controlled hypotension combined with the local use of adrenaline as an aid to the fenestration operation *MJ Australia* 1 300 1953
- 348 ORGANE G Decamethonium (bistrimethylammonium decane diiodide) in anaesthesia *Lancet* 1 773 1949
- 349 ——— Hypotension in anaesthesia *Anesth & Analg* 32 19 1953
- 350 ——— PATON W H M and ZAIMS E J Preliminary trials of bistrimethylammonium decane and pentane diiodide (C10 and C5) in man *Lancet* 1 21 1949
- 351 PAGE I H The occurrence of vasoconstrictor substance in blood during shock induced by trauma hemorrhage and burns *Am J Physiol* 139 386 1943
- 352 ——— Vascular mechanisms of terminal shock *Cleveland Clin Quart* 13 1 1946

- 353 ——— Treatment of shock by intra arterial infusion *Bull US Army M Dept* 7 366 1947
- 354 PAGLIARINI G Ipotensivi in chirurgia a scopo antiemorragico *Minerva farm Tor* 2 246 1953
- 355 PALLESTRINI E Lipotensione controllata per l'uso dei ganglioplegici nella chirurgia otorinolaringoatrica *Minerva otorinolaring*, 3 1 1953
- 356 PAPPER E M Renal function during general anesthesia and operation *JAMA* 152 1686 1953
- 357 PARKHOUSE J Cardiac effects of hypotension *Brit J Anaesth* 26 54 1954
- 358 PASQUALI E and CILLO L Ricerche sperimentali sulla funzionalità e l'istomorfologia del fegato nell'ipotensione da ganglioplegia *Acta chir patav* 9 117 1953
- 359 PATON W D M Paralysis of autonomic ganglia and the therapeutic effects of ganglion blocking drugs *Brit M J* 1 773 1951
- 360 ——— and THOMSON J W Procaine amide *Brit M J* 1 991 1953
- 361 ——— and WALKER J Methonium compounds in hypertension *Lancet* 1 473 1951
- 362 ——— and ZAIMIS E J Curare like action of polymethylene bis quaternary ammonium salts *Nature London* 161 718 1948
- 363 ——— and ZAIMIS E J Clinical potentialities of certain bis quaternary salts causing neuromuscular and ganglionic block *Nature London* 162 810 1948
- 364 ——— The pharmacological actions of polymethylene bistrimethylammonium salts *Brit J Pharmacol* 4 381 1949
- 365 ——— Paralysis of autonomic ganglia by methonium salts *Brit J Pharmacol* 6 155 1951
- 366 PATRONICOLAS P L'hypotension contrôlée par les sels de méthonium en chirurgie endothoracique *Praxis* 43 542 1954
- 367 PATURY E. and SOUZA A Hipotensao induzida em cirurgia geral *Rev brasil cir* 24 871 1952
- 368 PAYNE J P Controlled hypotension in theory and practice *Brit J Anaesth* 25 134 1953
- 369 ——— and BESWICK F B The oxygen consumption of the hypotensive cat *Brit J Anaesth* 26 252 1954
- 370 PAYNE R Subarachnoid injection of cocaine as a general anesthetic for operations upon the head *Tr Am Laryng Rhin & Otol Soc* 1 215 1902
- 371 PELMORE J F Death after anaesthetic with hypotension *Lancet* 2 1273 1952
- 372 PENAUD C L'hypotension contrôlée dans la prostatectomie *Bordeaux chir* 4 199 1953

- 373 PENN J., and BENTAL, H. Hypotensive anesthesia in plastic surgery *South African M J.*, 26:9 1952.
- 374 PENNY C. J., and SHACKLETON R. P. W.. A case of eclampsia treated with hexamethonium bromide. *Lancet* 2:617 1951
- 375 PETIT DUTAILLE D., and GUIOT G.. Indications et résultats de l'utilisation de certains ganglioplégiques (hypotension contrôlée) en chirurgie cérébrale. *J internat chir.*, 12:211 1952.
- 376 ———, ———. L'hypotension contrôlée en neuro-chirurgie *Mém Acad. chir.*, 79 373 1953
- 377 ———, ———. Indications et résultats du traitement chirurgical des anévrysmes orsoides et des anévrysmes artério-veineux du cerveau d'après 19 cas dont 14 traités par l'ablation de la lésion. *Presse méd.*, 61 1719 1953
- 378 ———, ———, and DAMONZEAU B.. De l'utilisation de certains ganglioplégiques hexaméthonium 92:95 C et des résultats qu'on peut obtenir en neuro-chirurgie. *Path gén* 52:651 1952.
- 379 PRIDEMSTER, D. B., LAESTER, C. H., EICHELBERGER, L., and SCHACHTER, R. J.. Afferent vasopressor nerve impulses as a cause of shock. tested experimentally by aortic-depressor nerve stimulation. *Ann Surg* 119:26 1944
- 380 PITKIN G. P.. Controllable spinal anesthesia. *Am. J Surg.*, 5 537 1928
- 381 POOLER, E. H.. Controlled hypotension in neurosurgery *Brit.M.J.*, 1 1163 1953
- 382 PORRO A., and AMORTH G.. Ipotensione controllata nella asportazione delle ernie del disco intervertebrale. *Minerva ortop.*, 4 421 1953
- 383 RAKITA L., and SANCIETTA S. M.. Acute hemodynamic effects of hexamethonium in normotensive man. *Circul Res.*, 1:499 1953
- 384 RANDALL, L. O., PETERSON W. G., and LERMAN G.. The ganglionic blocking action of thiophanium derivatives. *J Pharmacol & Exper Therap* 97 48 1949
- 385 REINHOLD H., and COERS C.. Controlled and uncontrolled hypotension. *Acta anaesth., belg.*, 3 18 1952.
- 386 RESTALL, P. A., and SMIRK, F. H.. The treatment of high blood pressure with hexamethonium iodide. *New Zealand M J* 49:206 1950
- 387 ———, ———. Regulation of blood pressure levels by hexamethonium bromide and mechanical devices. *Brit Heart J.*, 14 1 1952.
- 388 REYNOLDS T. B., PATON A., FREEMAN M., HOWARD F., and SHERLOCK, S.. The effect of hexamethonium bromide on splanchnic blood flow oxygen consumption and glucose output in man. *J Clin Investigation* 32:793 1953
- 389 RISKIN A. M., SEMERARO D., and ROBERTAZZI R. W.. The control

- 353 ——— Treatment of shock by intra arterial infusion *Bull US Army M Dept* 7 366 1947
- 354 PAGLIARINI G Ipotensivi in chirurgia a scopo antemorragico *Minerva farm Tor* 2 246 1953
- 355 PALLESTRINI E Lipotensione controllata per l'uso dei ganglioplegici nella chirurgia otorinolaringoatrica *Minerva otorinolaring* 3 1 1953
- 356 PAPPER E M Renal function during general anesthesia and operation *J.A.M.A* 152 1686 1953
- 357 PARKHOUSE J Cardiac effects of hypotension *Brit J Anaesth* 26 54 1954
- 358 PASQUALI E and CILLO L Ricerche sperimentali sulla funzionalità e istomorfologia del fegato nell'ipotensione da ganglioplegia *Acta chir patav* 9 117 1953
- 359 PATON W D M Paralysis of autonomic ganglia and the therapeutic effects of ganglion blocking drugs *Brit.M J* 1 775 1951
- 360 ——— and THOMPSON J W Procaine amide *Brit.M J* 1 991 1953
- 361 ——— and WALKER J Methonium compounds in hypertension *Lancet* 1 475 1951
- 362 ——— and ZAIMIS E J Curare like action of polymethylene bis quaternary ammonium salts *Nature London* 161 718 1948
- 363 ——— and ZAIMIS E J Clinical potentialities of certain bis quaternary salts causing neuromuscular and ganglionic block *Nature London* 162 810 1948
- 364 ——— The pharmacological actions of polymethylene bistrimethylammonium salts *Brit J Pharmacol* 4 381 1949
- 365 ——— Paralysis of autonomic ganglia by methonium salts *Brit J Pharmacol* 6 155 1951
- 366 PATRONICOLAS P L'hypotension contrôlée par les sels de méthonium en chirurgie endothoracique *Praxis* 43 542 1954
- 367 PATURY E. and SOUZA A Hipotensao induzida em cirurgia geral *Rev brasil cir* 24 871 1952
- 368 PAYNE J P Controlled hypotension in theory and practice *Brit J Anaesth* 25 134 1953
- 369 ——— and BESWICK F B The oxygen consumption of the hypotensive cat *Brit J Anaesth* 26 252 1954
- 370 PAYNE R Subarachnoid injection of cocaine as a general anesthetic for operations upon the head *Tr Am Laryng Rhin & Otol Soc* 1 215 1902
- 371 PELMORE J F Death after anaesthetic with hypotension *Lancet* 2 1275 1952
- 372 PENAUD C L'hypotension contrôlée dans la prostatectomie *Bardeux chir* 4 199 1953

- 373 PENN J., and BENTAL, H. Hypotensive anesthesia in plastic surgery
South African M J 26:9 1952
- 374 PENNY C. J., and SHACKLETON R. P. W., A case of eclampsia treated
with hexamethonium bromide *Lancet* 2:617 1951
- 375 PETIT DUTAILLIS D., and GUIOT G. Indications et résultats de l'utili-
sation de certains ganglioplégiques (hypotension contrôlée) en
chirurgie cérébrale. *J internat chir* 12:241 1952
- 376 ———, ———, L'hypotension contrôlée en neuro chirurgie
Mém Acad chir. 79 373 1953
- 377 ———, ———, Indications et résultats du traitement chirur-
gical des anévrysmes carotides et des anévrysmes artério-veineux du
cerveau d'après 19 cas dont 14 traités par l'ablation de la lésion
Presse méd. 61 1719 1953
- 378 ———, ———, and DAMONZEAU B., De l'utilisation de cer-
tains ganglioplégiques hexaméthonium 929: C et des résultats
qu'on peut obtenir en neuro-chirurgie *Path gén* 52:601 1952
- 379 PRENSTER, D. B., LAESTER C. H. EICHELBERGER, L. and SCHACHTER
R. J., Afferent vasopressor nerve impulses as a cause of shock
tested experimentally by aortic-depressor nerve stimulation *Ann
Surg* 119:26 1944
- 380 PITKEN E. P., Controllable spinal anesthesia *Am J Surg* 5:537
1928
- 381 POOLER E. H., Controlled hypotension in neurosurgery *Brit.M.J.*,
1 1163 1953
- 382 PORRO A., and AMORTH G., Ipotensione controllata nella asporta-
zione delle ernie del disco intervertebrale *Minerva ortop* 4 421
1953
- 383 RAKITA L. and SANCETTA S. M., Acute hemodynamic effects of
hexamethonium in normotensive man *Circul Res* 1 499 1953
- 384 RANDALL, L. O., PETERSON W. G., and LEHMANN G., The ganglionic
blocking action of thiophanium derivatives *J Pharmacol & Exper
Therap* 97:48 1949
- 385 REINHOLD H. and COERS C., Controlled and uncontrolled hypoten-
sion. *Acta anaesth., belg* 3 18 1952
- 386 RESTALL, P. A., and SMIRK F. H., The treatment of high blood pres-
sure with hexamethonium iodide *New Zealand M J* 49:206 1950
- 387 ———, ———, Regulation of blood pressure levels by hexa-
methonium bromide and mechanical devices. *Brit Heart J* 14 1
1952.
- 388 REYNOLDS T. B., PATON A. FREEMAN M. HOWARD F. and SHERLOCK
S., The effect of hexamethonium bromide on splanchnic blood
flow oxygen consumption and glucose output in man *J Clin
Investigation* 32 793 1953
- 389 RISKIN A. M., SEMERARO D., and ROBERTAZZI R. W., The control

- and management of hypertensive crises developing during surgical procedures *Anesthesiology* 15:262 1954
- 390 ROLLASON W N Anaesthesia and the reduction of blood loss *Anaesthesia* 7:10 1952
- 391 ———— Anaesthesia and the bloodless field. *Anesth & Analg* 32:289 1953
- 392 ROSENHEIM M L. Bromism on low-salt diets *Lancet* 1:347 1951
- 393 ROSENTHAL, A. Prostatektomie in kontrollierter blutdrucksenkung *Arch klin Chir* 276:316 1953
- 394 ROSENTHAL, A I Anesthesia for intracranial surgery *Am J Surg.* 85:673 1953
- 395 ROTTGEN P Über die operative behandlung sackförmiger hirnaneurysmen unter künstlicher blutdruckherabsetzung *Zentralbl Neuro chir* 12:200 1952
- 396 ROWLANDS E N WOLFF H. II and ATKINSON M Clinical assessment of drugs which inhibit gastric secretion with special reference to hexamethonium *Lancet* 2:1155 1952
- 397 RUTKOWSKI J, POKRZYWICKI S and HANKIEWICZ, J. Kontrolowane podciśnienie krwi *Polski tygodnik lek* 7:1358 1952
- 398 ———— ———— ———— Untersuchungen über die gesteuerte blutdrucksenkung *Schweiz. med Wchnschr* 83:212 1953
- 399 RYCROFT B W., and ROMANES G J Preliminary report on the use of methonium compounds and general anaesthesia in ophthalmic surgery *Brit J Ophth* 36:29 1952
- 400 SADOVE, M S WYANT G M GRITTELSON L. A., and BUCY P C. Controlled hypotension *J Neurosurg* 10:272 1953
- 401 SADOVE, M S WYANT G M and GLEAVE, G Controlled hypotension A study on arfonad (RO 2 2222) *Anaesthesia* 8:175 1953
- 402 ———— ———— ———— Thimethaphan camphorsulfonate (arfonad) in general surgery *A.M.A Arch Surg* 68:779 1954
- 403 ———— ———— ———— and BUCY P C. Controlled hypotension II A preliminary report on "arfonad" (RO 2 2222) *J Neurosurg.* 11:143 1954
- 404 SALGADO A. S and NIETO J C. Rol del anestesista en el control de la hemorragia quirúrgica *Rev argent anestesiol* 15:7 1953
- 405 SALLS S von Hypotonie bei prostatektomien *Helvet chir acta* 21:119 1954
- 406 SAMAIN A. La rachipertaine fractionnée comme anesthésie de base en chirurgie majeure son rôle dans la contrôle des médiateurs chimiques de l'influx nerveux la circulation contrôlée *Acta chir belg* 50:456 1951
- 407 ———— A propos de la chirurgie exsangue par circulation contrôlée. *Acta chir belg.* 50:634 1951
- 408 SANCTIYA, E M., LYNN R. B., SIMONEZ F A. HECKMAN G., and

- JANOUSKOVC, H.. Studies of hemodynamic changes in humans following induction of spinal anesthesia IV Observations in low spinal anesthesia during surgery *Surg Gynec & Obst* 97 597 1953
- 409 SANCETTA S M LYNN R B SIMONE F A and SCOTT R W Studies of hemodynamic changes in humans following induction of low and high spinal anesthesia I General considerations of the problem *Circulation* 6 559 1952
- 410 SARNOFF S J.. Massive pulmonary edema of central nervous system origin hemodynamic observations and the role of sympathetic pathways *Federation Proc*, 10 118 1951
- 411 ——— and ARRONOOD J G Differential spinal block A preliminary report. *Surgery* 20 150 1946
412. ———, ——— Differential spinal block II The reaction of sudomotor and vasomotor fibers *J Clin Investigation* 26 203 1947
- 413 ———, ——— Differential spinal block III The block of cutaneous and stretch reflexes in the presence of unimpaired position sense *J Neurophysiol* 10 205 1948
- 414 ———, ———, and CHAPMAN W P Differential spinal block IV The investigation of intestinal dyskinesia colonic atony and visceral afferent fibers *Surg Gynec & Obst* 86 571 1948
- 415 ——— and BERGLUND E Neurohemodynamics of pulmonary edema IV Effect of systemic vasoconstriction and subsequent vasodilatation on flow and pressures in systemic and pulmonary vascular beds *Am J Physiol* 170 588 1952
- 416 ———, ———, and SARNOFF L C Neurohemodynamics of pulmonary edema III Estimated changes in pulmonary blood volume accompanying systemic vasoconstriction and vasodilation *J Appl Physiol* 5 367 1953
- 417 ——— GOODALE W T and SARNOFF L C Graded reduction of arterial pressure in man by means of a thiophanium derivative (RO 11 2222) Preliminary observations on its effect in acute pulmonary edema *Circulation* 6 63 1952
- 418 ——— and SARNOFF L C Neurohemodynamics of pulmonary edema I Autonomic influence on pulmonary vascular pressures and the acute pulmonary edema state Preliminary report *Dis Chest* 22 685 1952
- 419 ———, ——— Neurohemodynamics of pulmonary edema II The role of sympathetic pathways in the elevation of pulmonary and systemic vascular pressures following the intracisternal injection of fibrin *Circulation* 6 51 1952
- 420 SAUNDERS J W Negative pressure device for controlled hypotension during surgical operations *Lancet* I 1286 1952

- 421 ———— Effect of controlled hypotension on cerebral function and circulation *Lancet* 1 1156 1954
- 422 SAUVAGE Y and HOFFMAN T. Erfahrungen und bemerkungen zu der kontrollierten blutdrucksenkung in der lungenchirurgie *Med Klin Berlin* 47 1515 1952
- 423 SAVILLE, S. Pentamethonium in hypertension *Lancet* 2 358 1950
- 424 SAVOLAINEN P V. Alhaisen verenpaineen käyttö leikkausvaurion hillitsemiseksi *Suom laak lehti* 24-993 1952
- 425 SCHALLEK W and WALZ, D. Effects of drug induced hypotension on the electro-encephalogram of the dog *Anesthesiology* 15-675 1954
- 426 SCHEINBERG P and STEAD E. A. JR. The cerebral blood flow in male subjects as measured by the nitrous oxide technique. Normal values for blood flow oxygen utilization and peripheral resistance with observations on the effect of tilting and anxiety *J Clin Investigation* 28 1163 1949
- 427 SCHNAPER, H W JOHNSON R. L. TLOHY E. B and FREIS E. D.. The effect of hexamethonium as compared to procaine or mety caine lumbar block on the blood flow to the foot of normal subjects *J Clin Investigation* 30 786 1951
- 428 SCHNEIDER, J. Über die anwendung des neuen ganglienblockierenden mittels pendiomid in der urologie. *Schweiz. med Wchnschr* 81 704 1951
- 429 ———— Pendiomid bei akuter lungenembolie *Helvet chir acta* 18 370 1951
- 430 SCHONBAUER, L. KUCHER R and STEINBEREITHNER, K. Prolongierte kontrollierte blutdrucksenkung ein beitrage zur konservativen behandlung intrakraneller blutungen und akuter hirndrucksteigerung *Wien med Wchnschr* 103 185 1953
- 431 SCHROEDER H A. The effect of 1 hydrazinophthalazine and hexamethonium on hypertension. *J Lab & Clin Med* 38-949 1951
- 432 SCHWEIZER, O. Clinical and investigational experiences with hypotension induced by hexamethonium bromide *New York J Med* 53 1951 1953
- 433 SCOTT L. D W KAY A W O'HARE, M M and SIMPSON J A. Hexamethonium bromide in duodenal ulcer *Brit.MJ* 2 1470 1950
- 434 SCULTETY S and JAKI J. Klinische und experimentelle beitrage zur künstlichen und regulierbaren hypotension *Zentralbl chir* 79 889 1954
- 435 SCURE, C. F.. Reduction of haemorrhage in the operative field by the use of pentamethonium iodide a preliminary report. *Anesthesiology* 12-253 1951
- 436 ———— Pentamethonium and shock *Lancet* 1 1020 1951
- 437 ———— and WYMAN J B. Controlled hypotension with arfonad. *Lancet* 1 338 1954

- 438 SEEVERS M H and WATERS R M Respiratory and circulatory changes during spinal anaesthesia *JAMA*, 99:961 1952
- 439 SCHACHTER M Hexamethonium and insulin hypoglycaemia *J Physiol* 113:206 1951
- 440 SHACKLETON R F W The reduction of surgical hemorrhage some observations on controlled hypotension with methonium compounds. *Brit.M.J.* 110:4 1951
- 441 ———, ———, The physiological background of controlled hypotension *Anaesthetist* 3:45 1951
- 442 SHACKMAN R and GRABER I G Reciprocal changes in blood flow of the limbs and rectum during general anaesthesia *Brit.M.J.* 11284 1952
- 443 ———, ———, and MELROSE D G The haemodynamics of the surgical patient under general anaesthesia *Brit. J. Surg.*, 40:193 1952
- 444 ———, ———, ——— Liver blood flow and general anaesthesia *Clin. Sc.* 12:307 1953
- 445 ———, ———, ——— and SMITH J Haemodynamics of methonium hypotension during anaesthesia *Anaesthesia* 7:217 1952
- 446 SHAPIRO A P and GROLLMAN A A critical evaluation of the hypotensive action of hydralazine hexamethonium tetraethyl ammonium and dibenzylamine salts in human and experimental hypertension *Circulation* 8:188 1953
- 447 SHORR, E. ZWEIFACH B W and FURCHGOTT R F On the occurrence sites and modes of origin and destruction of principles affecting the compensatory vascular mechanisms in experimental shock *Science* 102:489 1945
- 448 ———, ———, ——— Hepato renal factors in circulatory homeostasis III The influence of humoral factors of hepato renal origin on vascular reactions to hemorrhage *Ann. New York Acad. Sc.* 49:571 1948
- 449 SICARD A and SIBAUB Y L'hypotension contrôlée en chirurgie générale *Mém Acad. chir.* 79:361 1953
- 450 ———, ——— La chirurgie sous hypotension contrôlée *Rev. méd. Moyen Orient* 10:441 1953
- 451 SIGUIER F DAMOISEAU B and GUIOT G L'emploi des ganglioplégiques en neuro-chirurgie Avantages Technique Etude physiologique *Bull. et mém. Soc. méd. hôp. Paris* 67:1201 1951
- 452 SIMS J L MORRIS L E ORTH G S and WATERS R M The influence of oxygen and carbon dioxide levels during anesthesia upon post-surgical hepatic damage *J. Lab. & Clin. Med.* 38:388 1951
- 453 SMIRK F H Methonium compounds in hypertension *Lancet* 2:477 1950

- 454 ——— Practical details of the methonium treatment of high blood pressure *New Zealand M J* 49 637 1950
- 455 ——— Methonium compounds in high blood pressure *Lancet* I 346 1951
- 456 ——— and ALSTAD K E Treatment of hypertension by penta and hexa methonium salts *Brit.M J* I 1217 1951
- 457 SMITH H W *The Kidney Structure and Function in Health and Disease* New York Oxford University Press 1951
- 458 SNYDER H B MORRIS G C MOYER J H and HAYNES B W, JR. The effect of hexamethonium induced hypotension on water and electrolyte excretion in man *Surg Forum* 4 584 1953
- 459 SOFFER A and SWEET R H Effect of position changes of the lower extremities during vasomotor block *J.A.M.A* 151 1191 1953
- 460 SOMMERVILLE J and ALIAN J A case of drug eruption induced by hexamethonium bromide *Brit.M J* I 864 1951
- 461 STEPHEN C R WOODHALL B ODOM G NOWILL W K and BLOOR B Methods for reduction of surgical blood loss *J Kentucky M.A* 52 344 1954
- 462 STEVEN R J M and TOVELL, R M Hexamethonium salts as an aid to the orthopedic surgeon *J.A.M.A* 154 402 1954
- 463 STONE H H Controlled hypotension during surgery *Maryland M J* 3 163 1954
- 464 ——— MACKRELL T N and WECHSLER R L The effects of hypotension induced by intravenous hexamethonium and head up tilt upon cerebral hemodynamics in man *Am J M Sc* 228 112 1954
- 465 STOVNER J and LINDELL S Controlled hypotension to reduce bleeding in operative field *Nord med* 47 206 1952
- 466 SWANSON H S GALVIN W H and FINCHER E F Total spinal anesthesia in the surgical management of a congenital arteriovenous aneurysm *Surg Gynec & Obst* 93 215 1951
- 467 SWERDLOW M and WADE J W The effects of hexamethonium iodide on the electrocardiograph in anaesthetized subjects *Brit J Anaesth* 25 284 1953
- 468 SZUTU C and SCHWEIZER O Use of hexamethonium bromide for the experimental production of controlled hypotension in the anesthetized dog *Anesth & Analg* 33 18 1954
- 469 TAIT H and CAGLIERI G Experimental and clinical notes on the subarachnoid space *J.A.M.A* 35 6 1900
- 470 TESTA G Lipotensione provocata a scopo antiemorragico in chirurgia *Rassegna med sarda* 54 261 1952
- 471 TORRES S R Hipotensao controlada sau utilizacao em 32 casos cirurgicos *Med contemp* 71 231 1953

- 472 TURNER R Medical sympathectomy in hypertension a clinical study of methonium compounds *Lancet* 2 353 1950
- 473 ——— Methonium compounds in hypertension *Lancet*, 1408 1951
- 474 UHLEN A and LUND J S Continuous spinal anesthesia in intra cranial procedures *Proc Staff Meet Mayo Clin* 27 317 1952
- 475 ULLMANN T D and DIENCOFF D Effect of hexamethonium (CG) on renal hemodynamics in man *AMA Arch Int Med* 92:228 1953
- 476 VALLE G Le sostanze ad azione bloccante gangliare nel campo ostetrico e ginecologico I L'uso dei ganglioplegici nella medicazione preanestetica e nella ipotensione controllata *Minerva ginec*, 5:677 1953
- 477 VAN BERGEN F H BUCKLEY J J FRENCH L A DOBSON A B and BROWN I A Physiologic alterations associated with hexamethonium induced hypotension *Anesthesiology* 15 507 1954
- 478 VAN DE WALLE J Quelques considérations sur l'hypotension contrôlée pendant la narcose *Acta anaesth belg* 2 49 1951
- 479 VAN WIEN W Note préliminaire sur l'hypotension contrôlée *Acta anaesth. belg* 2 76 1951
- 480 ——— Faut il compenser ou prévenir les pertes sanguines en chirurgie thoracique *Acta anaesth belg* 4:22 1953
- 481 VEAL, J R RUSSELL A S and ASHBAURN F S The reaction of the coronary blood pressure to graded stages of shock (hemorrhage) *Ann Surg* 136 485 1952
- 482 VEHRIS G R Heart beat and respiration in total novocaine anesthesia *Northwest Med* 30:256 322 1931
- 483 VENTO RUIZ V Hipotensión controlada en neurocirugía *Rev españ oto neuro oftal* 13 32 1954
- 484 VERNEJOU R Hypotension contrôlée et réfrigération *Marseille méd* 90 420 1953
- 485 VETTER H GRABNER, G MŁCZOCZ F and STEINBEREITHNER K Herzkatheter untersuchungen bei massiver blutdrucksenkung durch hexamethonium *Klin Wchnschr* 32 97 1954
- 486 VOORHOEVE H C Hemodynamics of controlled hypotension during anesthesia *Internat J Anesth* 1 16 1953
- 487 ——— Kunstmatige bloeddrukverlaging bij gynaecologische operaties *Nederl tijdschr verlosk* 53 329 1953
- 488 ——— Haemodynamische problemen bij hypotensie *Nederl tijdschr geneesk* 98 595 1954
- 489 ——— and DELAHAYE R Un nouvel hypotenseur contrôlable le RO 2 2222 (arfonad) *Acta anaesth belg* 4 83 1953

- 454 ——— Practical details of the methonium treatment of high blood pressure *New Zealand M J* 49 637 1950
- 455 ——— Methonium compounds in high blood pressure *Lancet* 1 346 1951
- 456 ——— and ALSTAD K S Treatment of hypertension by penta and hexa methonium salts *Brit.M J* 1 1217 1951
- 457 SMITH H W *The Kidney Structure and Function in Health and Disease* New York Oxford University Press 1951
- 458 SNYDER H B MORRIS G C MOYER J H and HAYNES H W JR. The effect of hexamethonium induced hypotension on water and electrolyte excretion in man *Surg Forum* 4 584 1953
- 459 SOFFER A and SWEET R B Effect of position changes of the lower extremities during vasomotor block *J.A.M.A* 151 1191 1953
- 460 SOMMERVILLE J and ALLAN J A case of drug eruption induced by hexamethonium bromide *Brit.M J* 1 864 1951
- 461 STEPHEN C R WOODHALL B ODOM G NOWILL, W K and BLOOR H Methods for reduction of surgical blood loss *J Kentucky M.A* 52 344 1954
- 462 STEVEN R J M and TOVELL, R M Hexamethonium salts as an aid to the orthopedic surgeon *J.A.M.A* 154 402 1954
- 463 STONE H H Controlled hypotension during surgery *Maryland M J* 3 163 1954
- 464 ——— MACKRELL T N and WECHSLER R L The effects of hypotension induced by intravenous hexamethonium and head up tilt upon cerebral hemodynamics in man *Am J M Sc* 278 112 1954
- 465 STOVNER J and LINDELL S Controlled hypotension to reduce bleeding in operative field *Nord med* 47-206 1952
- 466 SWANSON H S GALVIN W H and FINCHER, E F Total spinal anesthesia in the surgical management of a congenital arterio venous aneurysm *Surg Gynec & Obst* 93-215 1951
- 467 SWERDLOW M and WADE J W The effects of hexamethonium iodide on the electrocardiograph in anaesthetized subjects *Brit J Anaesth* 25-284 1953
- 468 SZUTU C and SCHWEIZER O Use of hexamethonium bromide for the experimental production of controlled hypotension in the anesthetized dog *Anesth & Analg* 33 18 1954
- 469 TAIT H and CAGLIERI G Experimental and clinical notes on the subarachnoid space *J.A.M.A* 35 6 1900
- 470 TESTA G Lipotensione provocata a scopo antiemorragico in chirurgia *Rassegna med sarda* 54-261 1952
- 471 TORRES S R Hipotensao controlada sau utilizacao em 32 casos cirurgicos *Med contemp* 71 231 1953

- 507 WOODHALL, B ODOM G L STEPHEN C H McCURT C., and NEILL, W R. The use of chemically induced hypotension in the treatment of intracranial aneurysm *North Carolina M J* 14 513 1953
- 508 WOOLSEY R. D and FREEMAN H J Induced hypotension in surgery of intracranial vascular anomalies report of 2 cases. *J Missouri M A* 29 316 1952
- 509 WRIGHT A B Spinal analgesia with special reference to operations above the diaphragm *N Standard* 549 1931
- 510 WYMAN J E Cox R and GILLIES J Discussion on hypotension during anaesthesia *Proc Roy Soc Med* 46-605 1953
- 511 YONKMAN F F The pharmacology of newer agents employed in the treatment of gastrointestinal disturbances with special reference to regional ileitis and ulcerative colitis *Bull New York Acad Med* 28-700 1952
- 512 YOUNG I M DE WARDENER, H E. and MILES B E Mechanism of the renal excretion of methonium compounds *Brit M J* 2 1500 1951
- 513 ZAJNIS E. J The synthesis of methonium compounds their isolation from urine and their photometric determination *Brit J Pharmacol* 5 424 1950
- 514 ZIMMERMAN K Vorteile und gefahren der hypotensionsverfahren bei chirurgischen eingriffen *Helvet chir acta* 19-293 1952
- 515 ZUGLIANI J A Novos aspectos da neurocirurgia em face da hipotensão induzida por substâncias ganglioplégicas *Rev brasil cir* 25 322 1953
- 516 ZUKSCHWERDT L Operationen in blutleere durch ganglienblockade (pendiomid) und lagerung *Deutsche med Wchnschr* 77 460 1952
- 517 ZWEIFACH B W HERSHEY S G ROVENSTINE E A LEE R E and CHAMBERS R Anesthetic agents as factors in circulatory reactions induced by hemorrhage *Surgery* 18 48 1945

- 490 ——— and OKKER J J Onbloedig opereren door middel van hypotensie *Nederl tijdschr geneesk* 97 1348 1953
- 491 VOURGH G L hexamethonium en neurochirurgie *Anesth et analg* 9 376 1952
- 492 ——— Sur trois cas d'hypotension contrôlée par hexaméthonium en neurochirurgie *Gaz med France* 59 163 1952
- 493 ——— L hexaméthonium agent hypotenseur en chirurgie *Presse med* 60 159 1952
- 494 VYSLONZIL E Die kontrollierte blutdrucksenkung und ihre bedeutung für gehorverbessernde operationen (fenestrationen) *Wien klin Wchnschr* 64 772 1952
- 495 WALKER G LEVY L H HYMAN A and ROMNEY R Prolonged hypotensive reaction to hexamethonium *JAMA* 154 1079 1954
- 496 WATSCHINGER B KUCHER R STEINBEREITHNER K VYSLONZIL E and WIRTINGER W Künstliche blutdrucksenkung und nierenfunktion *Anaesthesist* 3 54 1954
- 497 WEBER H F Das ganglionblockierende pendiomid (ciba) bei urologischen erkrankungen und operationen *Wien med Wchnschr* 103 451 1953
- 498 WENGER R DONEFF D and VYSLONZIL E Zur kreislaufwirkung ganglienblockierender stoffe praoperative erkennung der durch eine kontrollierte blutdrucksenkung gefährdeten patienten *Deutsche med Wchnschr* 78 322 1953
- 499 WERLO L FRISK A R WADE G and ELIASCH H Effect of hexamethonium bromide in arterial hypertension *Lancet* 2 470 1951
- 500 WIGGERS C J Myocardial depression in shock a survey of cardiodynamic studies *Am Heart J* 33 633 1944
- 501 WIKLUND P E Controlled hypotension at intracranial operations *J Neurosurg* 10 617 1953
- 502 WILFLINGSIEDER P Die medikamentöse blutdrucksenkung zur verhütung der operationsblutung *Deutsche med Wchnschr* 77 619 1952
- 503 ——— Weitere beobachtungen und erfahrungen mit der medikamentösen ganglien blockierung während der operation *Chirurgia* 24 105 1953
- 504 WILKINS R W HALPERIN M H and LITTEY J Effect of dependent position upon blood flow in limb *Circulation* 2 373 1950
- 505 WILSON V H and KEELEY K J The haemodynamic effects of hexamethonium bromide in patients with pulmonary hypertension and heart failure *South African J M Sc* 18 125 1953
- 506 WIRTINGER W Erfahrungen mit der künstlichen blutdrucksenkung in der operativen hals nasen ohrenheilkunde *Anaesthesist* 2 1953

Author Index

A

Adnani J. 106
 Alexander G. L. 106
 Allan J. 51 152
 Alstad K. S. 50 132
 Alvisi C. 116
 Amorth G. 96 127
 Anderson C. D. 34 35 122
 Anderson S. 50 94 106
 Anduze H. 29 52 54 121
 Angles C. 52 95 111
 Anlyan W. G. 71 106
 Apgar V. 106
 Arden E. P. 17 29 123
 Arnold P. 50 66 106
 Arrowood J. G. 42 106 129
 Aserman D. 52 78 94 107
 Ashburn F. S. 133
 Asquith E. 52 107
 Assali V. S. 56 72 107 119
 Atkinson M. 128

B

Baars J. H. 107
 Bacher K. R. 60 107
 Ballantine R. I. W. 34 56 94 107
 Bandiera C. 107
 Banton A. H. 50 51 107
 Barbier P. 107
 Barlow R. B. 14 107
 Barnes M. 52 54 66 112
 Barnett A. J. 50 107
 Bassett D. L. 8 108
 Bataille L. 52 119 120
 Beaconsfield P. 107
 Beecher H. K. 8 45 105 107 108
 115
 Bein H. J. 60 99 108
 Belfrage D. 108

Belasso E. 116
 Benke A. 60 108
 Bennett H. S. 8, 108
 Bennett W. A. 108
 Benson W. R. 71 106
 Bentel H. 52 96 108 115 127
 Berglund E. 56 100 129
 Bernsmeier A. 60 108
 Bertolotto R. 108
 Beswick F. B. 65 126
 Biagini S. 116
 Biedermann G. 108
 Bier 41
 Bilslund W. L. 34 35 108
 Binet J. P. 123
 Bishop G. H. 42 117
 Bizard G. 108
 Blakeley C. G. 96 108
 Blalock A. 8 108
 Block M. A. 108
 Bloor B. 132
 Blum R. 109
 Bodman R. I. 109
 Bonnycastle D. D. 56 114
 Borri B. L. 34 109
 Boulton T. B. 58 107
 Bourne W. 41 109 116
 Bousquet E. 122
 Boyan C. P. 28 52 54 95 96 109
 Bozza M. 109
 Bradley S. E. 116-117
 Brown A. S. 109
 Brown I. A. 133
 Brown J. M. 74 110
 Bromage P. R. 46-47 67 70 71
 77 109
 Brunet D. Aubiac E. 95 109
 Brunswick A. 4 52 54 95 96
 109
 Buckley J. J. 133

Bucy F C 56 128
 Bullough J H 109
 Bunker J P 45 115
 Burn R A 77 95 110
 Burt C C 50 54 55 66 109
 Burstein C L 40 42 47 110

C

Cagliari G 41 132
 Camerini F 65 109
 Cameron A J 77 95 110
 Campan L 29 52 54 121
 Campbell A 50 52 77 110
 Carbonera D 56 110
 Carette L 110
 Carravetta M 52 110
 Chambers R 8 35 135
 Chapman W P 42 129
 Chateaufreynaud J 52 95 110 119
 Christensen J C 60 94 110
 Churchill Davidson H C 72 73
 110 111 123
 Caliberti B J 99 110
 Cillo L 126
 Cobb C C 98 110
 Coers C 80 127
 Cole W H J 52 95 125
 Costa J M 110
 Cotten M deV 74 110
 Co Tui 40 42 47 110
 Coulter R L 37 66 68 119
 Courtin R F 34 35 122
 Cox R 99 135
 Crehan J P 13 14 55 125
 Crile G W 9 99 110
 Crosetti J F 60 100 111
 Crumpton C W 65 67 111
 Culbertson J W 72 120
 Cusack A L J 50 52 111
 Cusack P H 50 52 111

D

Dale H H 63 111
 Damoiseau B 13 14 60 94 116
 127 131

Danzig L E 56 100 111
 Daumet P 52 95 111
 Dauri A 95 111
 Davies R M 111
 Davison M H A 4 13 14 25
 52 54 66 74 75 98 111 112
 de Clive Lowe S G 56 111
 Delahaye R 56 133
 Dewar H A 67 89 111
 de Wardener H E 51 72 73 110
 111 123 135
 Diengott H 72 133
 Dobkin A B 133
 Docal J C 60 111
 Donati G S 111
 Doneff D 134
 Douglas D M 115
 Douglass R A Jr 72 107
 Douthwaite A H 50 111
 Driver R L 65 112
 Drugemoller P 60 100 125
 Duboucher H 100 111
 Du Bouchet N 52 95 112
 du Cailar J 95 112
 Dundee J W 56 112

E

Ebbli C 46 112
 Eckenhoff J E 65 112
 Eddleman E E Jr 112
 Effler D B 116
 Eichelberger L 62 127
 Eliasch H 50 51 64 134
 Emmett J L 112
 Enderby G E H 13 14 25 26 28
 52 54 66 73 92 112 113
 Erlanger J 114
 Espinasse J 52 119 120
 Esser H 60 108
 Etsten H 113
 Evans H 73 113

F

Fabris F 124
 Fanjeux G 113

Fazekas J F 50 113
 Ferguson L K 42 113
 Feruglio F S 113
 Fibak J 113
 Fincher E F 93 132
 Fine J 95 113
 Finnerty F A Jr 50 51 95 113
 114
 Fitzpatrick H F 116 117
 Fitzpatrick L J 47 95 115
 Flossdorf T 60 100 125
 Foltz E L 65 112
 Fomon S J 72 107
 Ford R V 113
 Foster A D Jr 42 47 125
 Fournier J. 52 95 122
 Francis I L 56 112
 Frankel E 50 113
 Freeman M 66 127
 Freeman N E 113
 Freihent H J 93 135
 Freis E D 50 52 64 66 67 93
 113 114 130
 French L A 133
 Frey R 114
 Frisk A R 50 51 64 134
 Furchgott R F 71 131

G

Galvin W H 95 132
 Gardner W J 13 30 32 34 35
 95 114
 Gasser H S 114
 Genazzani E 114
 Gerlach H 00 93 114
 Gertner S H 56 114
 Gilh G 114
 Gillan J G 79 114
 Gillies J 5 7 8 13 14 25 27 38
 40 42-44 52 54 61 63 66 88
 94 96 99 112 114 115 135
 Gilmore H R 63 67 115
 Guttelson L A 56 128
 Glasser H 35 115
 Gleave G 56 64 128

Glenn W W L 93 115
 Gmuchtel W 115
 Goetz R H 50 66 106
 Coffen B S 47 93 115
 Goldstein J 99 110
 Goldsmith A J H 115
 Goldstone B 115
 Gomer A E 56 100 111
 Goodale W T 56 100 129
 Goodman C F 52 93 118
 Goodyer A V N 93 115
 Graber I C 61 66 71 90 131
 Grabner G 115 133
 Graham A J P 50 54 55 66 109
 115
 Greene N M 45 115
 Gregoire N 108
 Grieve W S M 51 122
 Griffiths H W C 14 27 38 40
 42 44 63 88 96 115
 Griffiths J A 92 115
 Grob D 50 66 116
 Grollman A 131
 Grosskreutz D C 8 23 56-57 121
 Groves L K 116
 Guglielmi F 64 63 109 116
 Guidetti B 116
 Guot G 13 14 60 94 95 116
 127 131
 Gurd F H 41 116

H

Habif D V 116 117
 Haid H 65 117
 Hafkenschel J H 65 112
 Hale D E 32 35 95 114 117
 Hall J 98 110
 Hall R A 98 110
 Halperin M H 134
 Hampton L J 18 23 54 56 57
 61 79 115 117 121
 Handley C A 72 124
 Hankiewicz J 95 128
 Harper J K 100 117
 Harris H E 95 117

Bucy P C 56 128
 Bullough J S 109
 Bunker J P 45 115
 Burn R A 77 95 110
 Burt C C 50 54 55 66 109
 Burstein C L 40 42 47 110

C

Cagliari G 41 132
 Camerini P 65 109
 Cameron A J 77 95 110
 Campan L 29 52 54 121
 Campbell A 50 52 77 110
 Carbonera D 56 110
 Carotte L 110
 Carravetta M 52 110
 Chambers R 8 35 135
 Chapman W P 42 129
 Chateaufreyaud J 52 95 110 119
 Christensen J C 60 94 110
 Churchill Davidson H C 72 75
 110 111 123
 Ciliberti B J 99 110
 Cillo L 126
 Cobb C C 98 110
 Coers C 80 127
 Cole W H J 52 95 125
 Costa J 96 110
 Cotten M deV 74 110
 Co Tui 40 42 47 110
 Coulter R L 37 66 68 119
 Courtin R F 34 35 122
 Cox R 99 135
 Crehan J P 13 14 55 125
 Crile G W 9 99 110
 Crosetti J P 60 100 111
 Crumpton C W 65 67 111
 Culbertson J W 72 120
 Cusack A L J 50 52 111
 Cusack P B 50 52 111

D

Dale H H 63 111
 Damoiseau B 13 14 60 94 116
 127 131

Danng L E 56 100 111
 Daumet P 52 95 111
 Dauri A 95 111
 Davies R M 111
 Davison M H A 4 13 14 25
 52 54 66 74 75 98 111 112
 de Clive Lowe S G 56 111
 Delahaye R 56 133
 Dewar H A 67 89 111
 de Wardener H E 51 72 73 110
 111 123 135
 Diengott D 72 133
 Dobkin A B 133
 Docal J C 60 111
 Donati G S 111
 Doneff D 134
 Douglas D M 115
 Douglass R A Jr 72 107
 Douthwaite A H 50 111
 Driver R L 65 112
 Drugemoller P 60 100 125
 Duboucher H 100 111
 Du Bouchet N 52 95 112
 du Cailar J 95 112
 Dundee J W 56 112

E

Ebbin C 46 112
 Eckenhoff J E 65 112
 Eddleman E E Jr 112
 Effer D B 116
 Eichelberger L 62 127
 Ehasch H 50 51 64 134
 Emmett J L 112
 Enderby G E H 13 14 25 26 28
 52 54 66 73 92 112 115
 Erlanger J 114
 Espinasse J 52 119 120
 Esser H 60 108
 Etsten B 8 113
 Evans B 73 113

F

Fabris F 124
 Fanjeux G 113

- Landis E M 38 121
 Lang H 121
 Langford H G 66 116
 Langston K W 121
 Lassner J 96 119 121
 Latscha H I 52 95 112
 Lawrence H R 121
 Lazorthes G 29 52 54 121
 Leathart G L 121
 Learmouth J 121
 Le Brigand J 52 121
 Ledru R 100 121
 Lee R E H 35 135
 Le Fillatre G 41 121
 Lehmann G 55 127
 Leigh M B 41 109
 LeMoynes J 95 121
 Leonhardt K O 121
 Leveen H H 71 121
 Levy L 51 134
 Lewis I 52 63 121
 Lindell S 52 132
 Ling A 34 95 114
 Litchfield J W 51 118
 Litter J 134
 Little D M Jr 18 23 54 56 57
 61 62 76 79 81 89 114 117
 121
 Locket S 51 73 122
 Loew F 122
 Longtin J 52 95 122
 Longtin L 52 95 122
 Lorenz W 60 108
 Lorhan P H 122
 Lottenbach K 60 122
 Love J G 34 35 122
 Lower W E H 99 110
 Lowrance P 116 117
 Lundy J S 34 35 94 122 133
 Lynn R H 63 66 71 72 122 125
 128 129

 M
 Mackey W A 52 123
 MacKinnon J 51, 72 123
 Mackrell T N 67 132
 Magill I W 56 123
 Mandow E A 17 29 123
 Mangiavacchi F 96 124
 Mann F E 108
 Mann C 123
 Marland T J 123
 Marxer A 123
 Mason A A 17 52 60 123
 Mathey J 123
 McClure C 62 135
 McCormick, V 122
 McCubbin J W 122
 McGill R J 50 122
 McKissock W 56 94 106
 McLaughlin C R 52 122
 McMichael J 65 67 70 115 123
 McNally P A 50 52 123
 McQueen E G 50 51 72 123
 Meier R 60 99 108
 Melrose D E 64 66 71 90 131
 Mendelsohn M L 72 123
 Menezes L C 94 123
 Mennuni R 119
 Merle D Aubigne R 123
 Méthot J P 122
 Metz D B 117
 Meyerovitz L 95 113
 Miescher K 123
 Mignault G 52 95 122
 Milbled G 108
 Miles H E 51 72 73 110 111
 123 135
 Miller M I 122
 Millett M 94 124
 Mills L E 72 124
 Milne I G 51 65 67 115 124
 Milzoch F 115 133
 Molfetta N 96 124
 Mondini P 124
 Montag H 98 124
 Moon V H 53 124
 Moore F T 111

Harvey A M 50 116
 Hay E B 117
 Haynes B W Jr 67 72 117
 124 132
 Hayward G W 88 92 117
 Heckman G 128
 Heinbecker P 42 117
 Hermann H 40 42 47 117
 Hershey S G 8 35 117 135
 Hewer A J II 94 115 117 118

Heyer H E 112
 Heymans C 17 19 118
 Hickam J B 118
 Hienert G 90 118
 Higgins T F 64 66 67 114
 Hill K R 52 95 118
 Hillis B R 50 118
 Himwich H E 8 118
 Hirson C 51 118
 Hoffman T 52 95 118 130
 Holle F 118
 Holling H E 118
 Holt M C 51 118
 Hopkin D A B 56 118
 Horatz K 118
 Horwitz O 125
 Howard F 66 127
 Howell G R 43 109
 Hueber E F 118
 Huet P C 95 118
 Huggins P A 72 124
 Hughes G 52 95 118
 Hunter A M 52 III 118
 Hyman A 51 134

I

Ing H R 14 107
 Inglis A N 41 109

J

Jackson I 34 56 94 107 119
 Jaki J 130
 James A 37 66 68 119
 Jenkin P II 52 95 119

Jenkins A R 67 89 111
 Johnson R L 50 51 64 66-67
 114 130
 Joly R 52 90 122
 Jones R R 119
 Jonnesco T 41 119
 Jungling O 119
 Just O 60 119

K

Kaplan S A 72 107 119
 Kasman L P 41 120
 Kattus A A Jr 66 118
 Kay A W 50 119 130
 Keeley K J 100 134
 Keller J W 45 115
 Kelley R T 64 66 67 114
 Kelly J C C 50 118
 Kelsall A R 51 118
 Kenny S 52 119
 Kern E 52 96 118 120 123
 Kerr W S 45 115
 Kety S S 67-68 120
 Kielbowicz D 113
 Kilduff C J 56 120
 Kilpatrick J A 50 51 120
 Kirkendall W M 72 120
 Kirsch J 60 120
 Kohlstadt K G 30 31 III 120
 Kolbl H 120
 Kopelman H 65 67 115
 Korkis F II 25 52 54 66 112
 120
 Koster H 41 120
 Kronen P II 74 110
 Kucher R 60 95 100 118 120
 130 134
 Kuhlmann J 120

L

Labayle J 95 118
 Laborit H 120
 Laester C H 62 127
 Laidlaw P P 63 111

- Robertazzi R W 99 127
 Robertson E 50 52 77 110
 Roger M 112
 Rollason W N 28 52 63 128
 Romanes G J 52 95 128
 Romero C 113
 Rommey R 51 134
 Rose J C 50 64 66 67 114
 Rosenheim M L 50 51 66 106 128
 Rosenthal A 96 128
 Rosenthal A I 54 91 128
 Roth C. E. 114
 Roth F 95 118
 Rottgen P 95 128
 Rovenstone E A 8 33 42 47 99 100 125 135
 Rowlands E N 128
 Rowlands S 125
 Rycroft B W 52 95 128
 Ruggiero W F 40 42 47 110
 Ruu P 113
 Russell A S 135
 Russo A 114
 Rutkowski J 95 128
- S
- Sadove M S 56 98 128
 Saegado A S 128
 Saexinger H 118
 Salem M 71 106
 Salis M von 128
 Samain A 60 128
 Sancetta S M 63 66 71 72 122 125 127 129
 Sarnoff L C 56 100 129
 Sarnoff S J 13 14 42 55 56 100 106 125 129
 Saunders J W 18 36 37 66 68 119 129 130
 Sauvage Y 52 95 111 130
 Saville S 50 130
 Savolainen F V 130
 Scarborough W R 66 116
 Schachter M 131
 Schachter R J 62 127
 Schaller W 67 80 130
 Schecter A. E. 113
 Scheinberg P 130
 Schmidt C F 67-68 120
 Schnaper H W 50 51 54 66 67 114 130
 Schneider J 60 100 130
 Schonbauer L 100 130
 Schroeder H A. 50 52 130
 Schwartz M 52 119 120
 Schweizer O 71 72 130 132
 Scott L. D W 50 130
 Scott R. W 63 66 71 72 122 129
 Scultéty S 130
 Scurr C F 25 52 54 56 66 99 112 123 130
 Sedzimir C H 56 112
 Semerato D 99 127
 Seror J 100 111
 Shackleton R F W 25 28 50 52 54 66 73 112 127 131
 Shackman R 64 66 71 90 122 131
 Shaffer S A 113
 Shapiro A P 131
 Sherlock S 66 127
 Shingleton W W 71 106
 Shorr E 71 131
 Sibaud Y 131
 Sicard A 131
 Siemons K. 108
 Siguer F 60 94 131
 Simeone F A 63 66 71 72 122 128 129
 Simpson J A 50 130
 Sims J L 131
 Smirk F H 37 50 52 120 127 131 132
 Smith A N 50 119
 Smith H W 132
 Smythe C M 116 117
 Snyder H B 67 72 117 124 132
 Soffer A 132
 Sommerville J H 132

Moricca G 56 124
 Morin G 40 42 47 117
 Morris G C Jr 67 72 117 124
 132
 Morris L E 131
 Morris N 50 124
 Mortimer P L F 34 35 124
 Morton A W 41 124
 Moyer J H 67 72 113 117 124
 132
 Mueller R P 66 71 125
 Muklerjee S R 125
 Mulder A G 71 121
 Muller C A 60 100 111
 Mundeleeer P 52 125
 Murphy E A 51 125
 Murphy Q R 65 67 111
 Musgrove A H 25 52 54 66 112

N

Natellis F 34 109
 Needleman N L 125
 Neill W R 62 135
 Neumann C 42 47 125
 Nicholson M J 13 14 55 125
 Nieto J C 128
 Nilsson E 67 69 125
 North J F 42 56 111 113
 Norpoth L 60 100 125
 Norvill W K 132

O

O'Brien D 52 95 125
 Odom G L 62 132 135
 O'Hare M M 50 130
 Okker J J 134
 O'Leary J L 42 117
 Oleesky S 51 124
 Organe G 49 51 52 125
 Orth O S 131
 Owen S G 67 89 111

P

Page I H 30 31 34 35 115 120
 122 125 126

Pagliari G 126
 Pallesstrini E 95 126
 Papper E M 106 116 117 126
 Parkhouse J 126
 Pasquali E 126
 Passaleco J 52 95 112
 Paton A 66 127
 Paton W D M 14 17 49 51
 125 126
 Patronicolas P 126
 Patury E 126
 Payne J P 26 55 126
 Payne R 41 126
 Peimore J F 17 52 54 60 77
 112 123 126
 Penaud C 126
 Penn J 52 96 108 127
 Penny C J 50 127
 Peruzzo L 111
 Peterson W G 55 127
 Petit Dutailh D 94 95 127
 Pettavel J 60 100 111
 Phemister D B 62 127
 Pitkin G P 41 127
 Pizzoli R 65 109 116
 Pokrzywnicki S 95 128
 Pooler E H 94 127
 Porro A 96 127
 Prokop F 71 121
 Proot R 52 125
 Pryor W W 118

R

Rakita L 127
 Randall L O 55 127
 Ranoverchi E 64 116
 Reinhold H 80 127
 Remouchamps L 52 125
 Resegotti E 56 110
 Restall P A 37 50 52 127
 Reynolds T B 66 127
 Ricci C 64 116
 Riskin A M 99 127
 Rizzi R 52 110
 Robelet A 108

Woodhall B 62 132 135

Woolsey R D 95 135

Wright A D 135

Wright Samson 19

Wyant G M 56 B1 128

Wylie W D 72 73 110 111 123

Wyman J B 25 52 54 56 66 99
112 123 130 135

Wynn Williams D 111

Y

Yonkman F F 56 135

York E E 8 113

Young I M 51, 135

Z

Zaimis E J 14 49 51 125 126
135

Zimmerman A. 135

Zugliani J A 94 135

Zukschwerdt L. 60 135

Zweifach B W 8 55 71 117 131
135

Sones F M Jr 116

Soulter J 113

Souza A 126

Spurr C L 113

Stacey R S 121

Stanton J R 114

Stead E A Jr 130

Steevers M H 131

Steinbereithner K 60 100 115
120 130 133 134

Stellet L 29 52 54 121

Stephen C R 59 62 71 106 132
135

Steven R J M 54 96 132

Stone H H 67 132

Stoneham F J R 17 29 123

Stovner J 52 132

Sturtzbecher F 118

Suriani U 94 124

Suyemoto R 56 107

Swain P G 51 122

Swanson H S 95 132

Sweet R B 132

Swerdlow M 65 66 132

Szutu C 72 123 132

T

Tada Y 72 107

Tait D 41 132

Taylor H C 71 106

Testa G 132

Thompson J W 17 126

Thorne M G 50 111

Todd D P 105 107

Torres R H 96 110

Torres S R 132

Tovell R M 54 96 132

Trautmann R 120

Tuohy E B 50 64 66 130

Turner R 50 133

U

Uihlein A 94 133

Ullmann T D 72 133

V

Valle G 133

Van Bergen F H 133

van de Walle J 60 133

Vanlerenberghe J 108

van Wien W 52 133

Vcal J R 133

Vehrs G R 41 133

Vento Ruiz V 133

Vernejoul R 99 133

Vetter H 115 133

Vial J 40 42 47 117

Vineberg A M 41 116

von Felsing J M 45 116

Voorhoeve H C 56 133 134

Vourch G 52 94 119 134

Vyslonzil E 52 95 134

W

Wade G 50 51 66 132 134

Wakim K G 108

Walker G 51 134

Walker J 51 126

Walz D 67 80 130

Waters R M 131

Watschinger B 134

Watson J 52 122

Webb J K 117

Weber H F 96 134

Wechsler R L 67 132

Wenger P 134

Werko L 50 51 134

Wiggers C J 35 134

Wiklund F E 94 134

Wilflingseder P 52 134

Wilkins R W 114 134

Willis K 112

Wilson V H 100 134

Wirtzinger W 95 131

Witkin L 50 113

Wohlraub K 118

Wolff H H 128

Wood P M 47 115

Subject Index

A

- Abdominal eviscerations 96
- Abdominal surgery 9,
 - position of patient 27 28
 - radical 95
 - spinal block 46
- Abdomino-perineal resection 96
 - position of patient 28
 - synchronous 82
- Acetylcholine 49 63
- Addison's Disease 91
- Adrenal pheochromocytoma re
 - removal of 23
- Adrenalectomy 96
- Age extremes of 89 91 92
- Air embolism 29 36 81
- Airway 10,
 - obstruction of 9
- Alpha rhythm 69 70
- Amino acids 71
- Ammonium salts 71
- Anemia III 91
- Anesthesia
 - administration of 9
 - chloroform 74
 - circulatory reflexes 8
 - conduction 22 70
 - cyclopropane 74
 - depth of and bleeding 8
 - friction within closed anesthetic circuit 9
 - general 70
 - deep 8 12 13 20
 - effect on bleeding during surgery 7 9
 - hypotensive 4
 - spinal III
 - trichlorethylene 74
 - vasodilatation 9
 - and venous return 25
- Anesthetic indications 98 99
- Aneurysms
 - congenital 33
 - ligation of 6
- Anhydrous cocaine 42
- Anociassociation 99
- Anoxia 42 67
 - cellular 69
 - cerebral 77 84
 - liver 70 71
 - myocardial 84
 - tissue 40
 - damage arteriotomy 36
- Anuria 77, 79 82
- Anxiety 92
- Aorta
 - atheroma 83
 - coarctation of 95
 - compression of 79
 - pressure decreased 35
- Aortic arch 26
- Aortic depressor nerve 62
- Aortic femoral thrombosis 79
- Apnea 10
- Arfonad 14 22 24 46 55 59 90
 - cardiac output 64
 - disadvantages 59
 - hemorrhage 59
 - oxygen consumption 65
 - urinary excretion 56
- Arm elevating and degree of hypotension 26
- Arrhythmias 74
- Arterial air embolism 81
- Arterial bleeding and reinfusion and shock syndrome 30
- Arterial compression 12
- Arterial pressure 13 16 17 62 72 98

Lempert fenestration operation
 3
 pressure 8
 and carbon dioxide absorp-
 tion 10
 arterial 66 69
 chloroform 8
 in deep general anesthesia 8
 level of 61 63
 local ■■■
 and height of operative site
 26
 readings
 arm cuff method 61
 Sanborn electromanometer
 61
 reduced 14 20 25 30 40 79
 103
 varying the level of 26
 vasopressor agents 73
 replacement of during surgery
 3
 vessels cut during surgery 5
 viscosity 17 89
 volume
 circulating 3 16 62
 reduction of 30 37
 control of in arteriotomy 35
 low 89 91
 total 103 105
 Bone grafts 96
 Bradycardia 89
 Brain
 edematous 97
 effect of controlled hypotension
 67 70
 electrical activity of ■■■
 ischemia to 28
 postural 69
 psychic strain 99
 swollen 97
 tension 35
 tumor
 surgery 30

 vascular 96
 Bromism 51
 Bromsulphalein retention 71

C

Cachectic patient 104
 Caesarean sections 94
 Cannula indwelling 104
 Capillary dilatation 63
 Capillary ooze 56
 control of extensive 96
 Carbon dioxide
 absorption and surgical bleed-
 ing 10
 retention 74
 tissues containing increased con-
 centration of 74
 Carcinoma 82 83 85
 Cardiac arrest 78 79 81 84
 Cardiac arrhythmias 74
 Cardiac disease 89 90
 Cardiac failure 81 84 89 90 100
 Cardiac index 64
 Cardiac massage 84
 Cardiac output 16 64
 factors controlling 17
 in deep general anesthesia 8
 methonium compounds 64
 position of patient 64
 Cardiac vessel surgery 98
 Cardio aortic sinus 26
 Cardio pulmonary disease 6
 Cardiovascular system 63 88
 collapse 78 79 81 84 85
 disease 92
 surgery 95
 Carotid sinuses 26
 Cautery 11
 Cellular intoxication 20
 profound generalized 9
 Cellular respiration 104
 Central nervous system depression
 99
 Cerebral anoxia 77 78 84

- control of in arteriotomy 34 35
 - decreased 18
 - effect of posture 26
 - maintaining normal 16
 - pulmonary 64
 - systemic 39
 - Arterial tone regulation of 19
 - Arteries elasticity of 17
 - Arteriolar bed dilated by sympathetic blockade 73
 - Arteriolar dilatation 14 62 63 73
 - Arterioles contraction and tone of 20
 - Arteriosclerosis 88
 - Arteriotomy 13 18 24 31 37 III
 - advantages of 34 35
 - air embolism 36
 - anoxic tissue damage 36
 - apparatus for 32
 - circulatory depression 35
 - clot formation 36
 - craniotomy 33 34
 - disadvantages 35 36
 - equipment 36
 - fenestration operation 34
 - hemorrhagic shock 35
 - heparin administration of 34
 - hepatic resections 34
 - infection 36
 - intracranial surgery 35
 - neurosurgical procedures 34
 - open heart operations 34
 - reinfusion 35
 - vasoconstriction 34 35
 - Arteriovenous oxygen difference 67 68
 - Artificial hibernation 99
 - Atheroma 83 84
 - Auricular pressure right measurement of 64
 - Autonomic activity effect of barbiturates 8
 - Autonomic ganglia
 - blocking agents 48
 - inhibition of synaptic transmission of impulses through 60
 - Awakening time 76 77
- B**
- Bainbridge reflex 63
 - Ballistocardiography 64
 - Barbiturates 51
 - autonomic activity 8
 - effect on hypothalamic centers 8
 - hypotension 8
 - respiratory depressant effects of 8
 - Bleeding
 - and depth of anesthesia 8
 - cerebrovascular 100
 - control of 11 15 103
 - during surgery 3 5 10
 - anesthetic factors 7 10
 - control of 11 15 103
 - factors influencing 6
 - extensive 104
 - prevention of 8 9
 - general anesthetic agents 7
 - and administration of 9
 - in wound 25 26
 - Blindness unilateral 79
 - Blood
 - clotting of
 - deficiencies in 6
 - during nitrous oxide anesthesia 7
 - general anesthetic agents 7
 - flow
 - hepatic 71
 - increase 50
 - mean cerebral 67 68
 - nitrous oxide technique for measuring 67
 - peripheral 66
 - splanchnic 66
 - hepatic 70 71
 - intra arterial infusion of 30

- Lempert fenestration operation
 3
 pressure 8
 and carbon dioxide absorp-
 tion 10
 arterial 66 69
 chloroform 8
 in deep general anesthesia 8
 level of 61 63
 local 28
 and height of operative site
 26
 readings
 arm cuff method 61
 Sanborn electromanometer
 61
 reduced 14 20 25 30 40 79
 103
 varying the level of 26
 vasopressor agents 73
 replacement of during surgery
 3
 vessels cut during surgery 5
 viscosity 17 89
 volume
 circulating 3 16 62
 reduction of 30 37
 control of in arteriotomy 35
 low 89 91
 total 103 105
- Bone grafts 96
 Bradycardia 63
 Brain
 edematous 97
 effect of controlled hypotension
 67 70
 electrical activity of 69
 ischemia 28
 postural 69
 psychic strain 99
 swollen 97
 tension 35
 tumor
 surgery 30
- vascular 96
 Bromism 51
 Bromsulphalein retention 71
- C
- Cachectic patient 104
 Caesarean sections 94
 Cannula indwelling 104
 Capillary dilatation 63
 Capillary ooze 86
 control of extensive 96
 Carbon dioxide
 absorption and surgical bleed-
 ing 10
 retention 74
 tissues containing increased con-
 centration of 74
 Carcinoma 82 83 85
 Cardiac arrest 78 79 81 84
 Cardiac arrhythmias 74
 Cardiac disease 89 90
 Cardiac failure 81 84 89 90 100
 Cardiac index 64
 Cardiac massage 84
 Cardiac output 16 64
 factors controlling 17
 in deep general anesthesia 8
 methonium compounds 64
 position of patient 64
 Cardiac vessel surgery 98
 Cardio aortic sinus 26
 Cardio pulmonary disease 8
 Cardiovascular system 88 88
 collapse 78 79 81 84 85
 disease 92
 surgery 95
 Carotid sinuses 26
 Cautery 11
 Cellular intoxication 20
 profound generalized 9
 Cellular respiration 104
 Central nervous system depression
 99
 Cerebral anoxia 77 78 84

- Cerebral : arterial venous oxygen difference 88
- Cerebral blood flow 89 90
- Cerebral circulation 83
- Cerebral function measurement of 67
- Cerebral hemodynamics 67 69
during induced hypotension 67
- Cerebral ischemia 88
- Cerebral metabolism 49 67 70
- Cerebral oxygen consumption 67
- Cerebral thrombosis 78 79 89
and anoxia 81 82
- Cerebral vascular resistance 67 89
- Cerebral vessels passive congestion of 29
- Cerebrovascular bleeding 100
- Cerebrovascular disease 88
- Chemical agents of escharotic nature 11
- Chemical ganglionic blockade 45
- Chest surgery 95
spinal analgesia 41
- Chloride preparations 51
- Chloroform 12 13
and coagulation 7
blood pressure 8
hypotension 8
- Chlorpromazine 11
- Circulating blood volume reduction of 30 37
- Circulation
complications to 78
continued 29
obstruction to 78
time 66 67
- Circulatory depression arteriotomy 35
- Circulatory reflexes
effect of muscle relaxants 9
in light planes of anesthesia 8
in deep general anesthesia 8
receptors in aortic arch 26
receptors in cardio-aortic sinus 26
- receptors in carotid sinuses 26
- Coagulation non anesthetic factors to prolong 5 7
- Cocaine 42
- Cold local application of and hemostasis 11
- Colitis ulcerative 56
- Coma 70
- Complications 75 80
blurred vision 77
cardiac arrest 78
cardiovascular collapse 78
involving kidneys 77 78
prolonged and delayed awakening time 76 77
rate 75
reactionary hemorrhage 76
thrombotic phenomena 78
- Conduction anesthesia 22 38-47
- Constrictor tone
interrupted 22
reduction of 7
- Contraindications 88 93
- Convulsions 68 70
- Coronary arteries 83
- Coronary disease 88
- Coronary flow 65
- Coronary insufficiency 66
- Coronary thrombosis 78 79 88
- Coronary vascular resistance 65
- Cortex electrical energy of 89
- Cortical rhythm profound disturbances of 70
- Cranial surgery partial sympathetic blockade 47
- Craniotomy 27 51 97
arteriotomy 33 34
- Cyanosis 85
- Cyclopropane 52
and coagulation 7
- Cystoscopy bladder tenesmus during 60

D

Dale's postural ischemia 15

- Decamethonium 49 51
 and muscular relaxation 9
 Déconnexion 99
 Decortication 95
 Defibrillation 81
 Dehydrogenated ergot prepara-
 tions 23
 Delta waves 69 70
 Dental extractions 94
 Diabetes mellitus 92
 Diaphragm moved downward 10
 Diastole prolonged 83
 Dibenamine 22
 Differential spinal block 42
 Duodide 22
 Diseases
 cardio pulmonary 6
 producing high degree of tox-
 icity 5
 d tubocurarine and muscular re-
 laxation 9
 Duodenal ulcer 50

E

- Eclampsia
 hypertension in 46
 pendiomide 60
 Edema acute pulmonary 56
 Efferent sympathetic outflow pa-
 ralysis of 14
 Electrical activity of brain 69
 Electrocardiography 65
 Electroencephalogram 69
 tracings 77
 Elimination of drugs from body
 77
 Emboli 67
 Endotracheal cuff inflated 79
 Ephedrine 74
 Epidural analgesia 22 40
 Epidural block 46-47 71
 disadvantages 47
 Esmarch bandages 11
 Esophago gastrectomy 95

- Etamon intravenous administra-
 tion of 65
 Ether 52
 and coagulation 7
 Ethylene and coagulation 7
 Extradural analgesia 40
 Extrasystoles 65
 Extremities
 lower
 application of negative pres-
 sure to 37
 application of vacuum appa-
 ratus 18
 elevated skin temperatures in
 50
 surgery upon 11
 Eyeball pressure on 79

F

- Femoral artery cannulating 30
 Fenestration operation 31
 arteriotomy 11
 partial sympathetic blockade 47
 position of patient 27
 Fibrinogen deficiency 5
 Flicker fusion frequency test 69
 Fowler's position 104
 semi 27

G

- Ganglionic blockade 18 28
 and transmission of noxious im-
 pulses 99
 cardiac output 64
 mean blood flow 86
 reduction of peripheral resist-
 ance 48 60
 synergistic effect with general
 anesthesia 77
 Ganglionic blocking agents 64
 easily injectable 14
 medical therapies 99
 therapy of cardiac failures 90
 Ganglionic synapses 9
 Gastrectomy 96

- Cerebral anterior venous oxygen difference 68
- Cerebral blood flow 89 90
- Cerebral circulation 83
- Cerebral function measurement of 67
- Cerebral hemodynamics 67 69
during induced hypotension 67
- Cerebral ischemia 68
- Cerebral metabolism 29 67 70
- Cerebral oxygen consumption 67
- Cerebral thrombosis 78 79 89
and anoxia 81 83
- Cerebral vascular resistance 67 89
- Cerebral vessels passive congestion of 29
- Cerebrovascular bleeding 100
- Cerebrovascular disease 83
- Chemical agents of escharotic nature 11
- Chemical ganglionic blockade 45
- Chest surgery 95
spinal analgesia 41
- Chloride preparations 51
- Chloroform 12 13
and coagulation 7
blood pressure 8
hypotension 8
- Chlorpromazine 11
- Circulating blood volume reduction of 30 37
- Circulation
complications to 78
continued 29
obstruction to 78
time 66 67
- Circulatory depression arteriotomy 35
- Circulatory reflexes
effect of muscle relaxants 9
in light planes of anesthesia 8
in deep general anesthesia 8
receptors in aortic arch 26
receptors in cardio-aortic sinus 26
- receptors in carotid sinuses 26
- Coagulation non anesthetic factors to prolong 5 7
- Cocaine 42
- Cold local application of and hemostasis 11
- Colitis ulcerative 56
- Coma 70
- Complications 75 80
blurred vision 77
cardiac arrest 78
cardiovascular collapse 78
involving kidneys 77 78
prolonged and delayed awaking time 76-77
rate 75
reactionary hemorrhage 76
thrombotic phenomena 78
- Conduction anesthesia 22 38-47
- Constrictor tone
interrupted 22
reduction of 7
- Contraindications 88 93
- Convulsions 68 70
- Coronary arteries 88
- Coronary disease 88
- Coronary flow 65
- Coronary insufficiency 66
- Coronary thrombosis 78 79 88
- Coronary vascular resistance 88
- Cortex electrical energy of 69
- Cortical rhythm profound disturbances of 70
- Cranial surgery partial sympathetic blockade 47
- Craniotomy 27 31 97
arteriotomy 33 34
- Cyanosis 85
- Cyclopropane 52
and coagulation 7
- Cystoscopy bladder tenesmus during 60

D

Dales postural ischemia 15

controlled 4 15 15 25
 advantages of arteriotomy 34
 85
 and cerebral metabolism 29
 anesthetic indications 98 99
 clinical practice of 23 24
 complications 75 87
 conduction anesthesia 38 47
 contraindications 88 93
 controlling degree of 26
 electrocardiographic tracings
 65
 epidural block 46
 ganglionic block 48
 history of 18
 indications 94 100
 medical 99 100
 modified technique of 56 58
 patients resistant to metho-
 nium compounds 37
 physiological considerations
 61 74
 position of patient 26 28
 post operative care 105
 reduction of peripheral resist-
 ance 26
 requirements 101 105
 selection of patients 101 102
 surgical indications 96 98
 techniques employed 23
 types of operative procedures
 95
 degree of 79 80
 duration of 80
 during complete sympathetic
 blockade 29
 hypovolemic 35 103
 induced 4 16 24
 arfonad 55
 epidural block 46
 physiological basis of tech-
 niques for 18
 intentional production of 3
 methonium induced 37
 normovolemic, 40 103

 partial sympathetic blockade 47
 persistent 79 91
 posture of patient 61
 reduction of peripheral resist-
 ance 38-47
 uncontrolled 80
 Hydralazine 20
 Hyperextension 27 28
 Hypertension 50 89
 control of 20
 malignant 90
 methonium compounds 82
 renal 65
 severe 90
 Hypnosis 20 43
 Hypoglycemia 92
 Hypotensive anesthesia 4
 Hypothalamic centers inhibition of
 7
 Hypothermia 11 99
 Hypovolemic hypotension 35
 Hypoxia 71
 Hysterectomy radical 96

I

Insulin 92
 Intestinal capillaries oxygen de-
 saturation in 70
 Intestinal peristalsis inhibition of
 50
 Intra arterial infusion of blood or
 plasma 80
 Intracranial aneurysms 96
 ligation of 94 95
 Intracranial interventions 6
 Intracranial surgery arteriotomy
 35
 Intracranial tension 89
 Intrapleural negative pressure 9
 Intrapulmonary pressure 9
 Intra thoracic surgery 6 25
 position of patient 27
 Intravascular pressure increased
 54
 Intravascular tension 98

- Gastric motility reduction of 50
 Gastric secretion inhibition of 50
 Gastric ulcer 50
 Glaucoma 95 98
 pendiomide 60
 Glomerular filtration 51 72 90
 Glomeruli hydrostatic pressure in 72
 Goiter toxic subtotal thyroidectomy 84
 Gravitational venous drainage 12
 Gravity
 above heart level 26
 below heart level 26
 Gynecological surgery 6
- H**
- Hand elevating and degree of hypotension 26
 Head
 and neck surgery 95
 head down position 28 29
 head up position 27 29
 operations on spinal analgesia 41
 Headache 46-47
 spinal analgesia 46
 Heart
 and increased bleeding 5
 contractile force of 74
 force of beat 17
 gravity above level of 26
 gravity below level of 26
 operations on open arteriotomy 34
 rate 17 63
 stroke volume 64
 Heat local application of and hemostasis 11
 Hematocrit 65
 determinations of 105
 Hemipelvectomy 85 86
 Hemoglobinuric nephrosis 86
 Hemorrhage 7 59 63 82 85 89 91 94 98 104
 during surgery 3
 control of gross 3 96
 general anesthetic agents 7 8
 reactionary 11 73 76 79 81 85 86
 reduction of pressure by 62
 replacement of blood 3
 severe 100
 Hemorrhagic shock 11
 arteriotomy 35
 Hemostasis 31
 position of patient, 12
 Hemostatic sponges 11
 Hepatic artery 70
 Hepatic disease 89 91 102
 Hepatic necrosis acute 90
 Hepatic resections major arteriotomy 34
 Hepatic vein oxygen content 71
 Herpes zoster 100
 pendiomide 60
 Hexamethonium 14 48 50 52 54 68 73 84
 antidote to decamethonium 51
 bromide 82
 coronary blood flow 55
 effect on brain 67-68
 flicker fusion frequency test 60
 halides 22
 hypotension anesthesia record 53
 intravenous 65
 iodide 83 85
 oxygen consumption 65
 peripheral blood flow 66
 splanchnic blood flow 66
 Hiccup intractable 50
 Hip operations on 11
 Histamine 63
 Homologous serum jaundice 35
 Hypotension
 abruptness of onset 80
 arfonad 55 57
 barbiturates 8
 chloroform 8

- patients resistant to 54
- position of patient 54
- post-operative care 54 55
- renal excretion of 51
- inulin clearance ratios 51
- Mid abdominal surgery spinal
 - block 46
- Mortalities 80-87
 - rate of 80 81
 - renal failure and anuria 81 82
- Motor blockade 14
- Motor paralysis 42
- Muscle
 - limb contraction of 25 26
 - paralysis 25
 - relaxants 10
 - relaxation 12 99
 - during surgery 9
 - skeletal during clinical anes-
thesia 49
 - vascular 6
- Myocardial anoxia 84
- Myocardial ischemia 11 66
- Myocardium weakened 74
- Myoneural junction 49

N

- Narcotic agents 10
- Nausea 11
- Neck dissection radical 27
- Necrosis
 - acute hepatic 90
 - ischemic 86
 - liver 71
 - pressure 79
- Negative pressure device 36
- Nephrectomy 79 96
- Nephrosis hemoglobinuric 86
- Nerve block
 - concentrations of procaine re-
quired 43
 - direct 11
- Neurosurgical procedures 94
 - arteriotomy 54
 - operative trauma 81 87

- position of patient 27
- retractors employed 78
- withdrawal of blood prior to
30 81
- Nitrates 23
- Nitrites 13
- Nitrous oxide 44 52
 - and coagulation 7
 - technique for measuring blood
flow 67
- Node dissection radical 97
- Non rebreathing techniques 10
- Nor adrenaline 71
- Noxious impulses 10
- Nupercaine 44

O

- Olfactory groove meningioma ex-
cision of 31
- Oliguria 77 79
- Open hip operation 96
- Operative site
 - elevation of 13 29
 - nature of 6
 - posture of patient 6
 - visibility in 97
- Ophthalmic surgery 95
- Organ tension 98
- Orthopedic surgery 96
- Otorhinolaryngological procedures
95
- Over heparinization 11
- Oxygen consumption 65 66
- Oxygen desaturation in intestinal
capillaries 70
- Oxygenation 104 105

P

- Palpitation 92
- Parasympathetic blockade 64
- Parasympathetic ganglia 48
- Paravertebral block 40
- Paravertebral sympathetic ganglia
blockade 22
- Partial sympathetic blockade 47

- Intravascular thrombosis 89
 - Intravenous barbiturates and coagulation 7
 - Intubation 52
 - Inulin clearance of 51
 - rate 72
 - Iodism 51
 - Irreversibility 35
 - Ischemia
 - cerebral 68
 - myocardial 35 66
 - postural 25 29
 - of brain 69
 - to brain 28
 - vital organs 27 28
 - Ischemic necrosis 86
 - Ischemic operative field 9
- J**
- Jack knife position 27
 - Jaundice homologous serum 35
- K**
- Kidney
 - complications involving 77 78
 - disease 51 90
 - effects on 71 73
 - excretion of arfonad 90
 - excretion of methonium compounds 90
 - morphological changes 72
 - position of patient 28
- L**
- Laminectomy ■ 96
 - Laparotomy 71
 - Laryngeal mucosa pressure necrosis of 79
 - Laryngectomy 95
 - Lateral jack knife position 28 29
 - Left ventricle calculated work of 65
 - Leg
 - elevating and degree of hypotension 26
 - pneumatic suction to 68
 - Lempert fenestration operation 3
 - 95 97
 - Ligation temporary 12
 - Lithotomy position 28 29
 - Liver
 - anoxia 70 71
 - blood supply to 70
 - effect on 70 71
 - failure 90
 - level of hypotension 70
 - necrosis 71
 - occult damage 71
 - oxygen supply 70
 - Lobectomy 95
 - Lower extremities operations on
 - position of patient 28
 - Lumbar disc prolapsed ■ 87
 - Lumen 83
 - Lungs and increased bleeding 5
- M**
- Mandibular resection 27
 - Mastectomy 6
 - radical 27 57 83 95
 - Mastoidectomy spinal analgesia 41
 - Medical indications 99 100
 - Medullary center muscles of 14
 - Meningiomas removal of 95
 - Meningismus 47
 - Mesenteric thromboses 78
 - Metabolic rate
 - increased basal 5
 - lowered 77
 - Metabolism cerebral 29
 - Methedrine 45 71 82 84
 - Methonium 90
 - compounds 14 24 48 55 66 91
 - administration of 52 54
 - anesthesia 52
 - blurred vision 77
 - cardiac output 64
 - dosage 51 53
 - effect on liver 71
 - medical therapy of hypertension 52

hypertension in 46
 Pressor reflexes in deep general anesthesia 8
 Pressure necrosis 79
 Priscol 22
 Procaine 42
 concentrations required for nerve block 43
 preganglionic sympathetic fibers 43
 subdural injection of 63
 Prone jack knife position 29
 Prostatectomy 96
 Proteinuria 73
 Prothrombin deficiency 5
 Pulmonary edema 81 86, 100
 Pulmonary embolism 100
 acute 60
 Pulmonary infarcts 81
 Pulse rate 63
 vacillating 65
 Pupils eye dilatation of 50
 Pyrexia 56

R

Radical hysterectomy 96
 Radical mastectomy 95
 position of patient 27
 Radical neck dissection 27 95
 Radical node dissection 97
 Radical pelvic surgery 28
 Reactionary hemorrhage 79 81
 85 88
 Red cell mass 103
 Regime 22
 Reinfusion arteriotomy 35
 Renal blood flow 90
 Renal damage 72 73
 Renal disease 89 91 102
 Renal epithelium 72
 Renal excretion of methonium compounds 51
 Renal failure and anuria 81 82
 Renal function
 complications involving 77 78

depression of 71
 Impaired 51
 Requirements 101 105
 Respiration
 assisted 9
 cellular 36
 level of pressure 40
 controlled 9
 depression 74
 effect of barbiturates 8
 failure 42
 increased resistance to 9
 muscles of 14
 Restlessness 92
 Retinal artery occlusion 79
 Retinal thrombosis 78 79
 Retractor anemia 78
 Retransfusion 18

S

Salivary secretion inhibition of 50
 Sanborn electromanometer 81
 Sclerosis vascular 51
 Semi Fowler's position 27
 Sensory blockade 14
 Sepsis spinal analgesia 46
 Shock 35 88 91
 chronic 104
 irreversible 80 103 104
 surgical 99
 syndrome method for studying 30
 treatment of 30
 Shoulder joints operations on 6
 Side tilt of operating table 27
 Skin
 cold clammy 88
 flaps 96
 grafts 96
 Somnolence 68
 Speech slurring 68
 Spinal analgesia 40 72
 high 63 64 71 74
 arteriovenous oxygen 65
 oxygen consumption 65

- Patent ductus arteriosus 95
- Pelvic evisceration 96
- Pelvic floor repair of position of patient 28
- Pelvic node dissection 96
- Pelvic surgery 96
 - radical position of patient 28
- Pendiomide 14 22 24 58 60
 - administration of 60
 - advantages 60
 - disadvantages 60
 - dosage 60
- Pentamethonium 14 48 50 70 82
 - antidote to decamethonium 51
 - halides 22
 - intravenous 65
 - peripheral blood flow 66
- Pentothal 43
 - intravenous 52
- Perinephritis experimental 65
- Peripheral circulation 83
- Peripheral resistance 16 18 20 38
 - reduction of 26 38-47
 - by interruption of sympathetic vasoconstrictor impulses 21
 - ganglionic blockade 48 60
- Peripheral respiratory paralysis
 - spinal analgesia 42
- Peripheral vascular disease 50
- Perspiration 68
- Pheochromocytoma removal of 98
- Phlebotomy 90
- Physiological considerations 61 74
- Plasma intra arterial infusion of 30
- Plastic surgery 96
- Platelet deficiency 5
- Pneumatic suction to legs 68
- Pneumonectomy 95
- Polycythemia 89
 - vera 91
- Porta caval shunt 96
- Portal vein 70
- Position of patient 6 104
 - abdominal surgery 28
 - abdomino perineal resection 28
 - cardiac output 64
 - craniotomies 27
 - during and after complete sympathetic blockade 29
 - effect on arterial pressure 26
 - head down 29
 - tilt 28
 - head up 29
 - tilt 27
 - intra thoracic operations 27
 - ischemia
 - of brain 69
 - to vital organs 27
 - jack knife 27
 - kidney operations 28
 - lateral jack knife 28 29
 - lithotomy 28 29
 - lower extremities operations on 28
 - methonium compounds 54
 - neurosurgical procedures 27
 - post operative 105
 - posterior fossa explorations 27
 - prone jack knife 29
 - radical mastectomy 27
 - radical pelvic surgery 28
 - repair of pelvic floor 28
 - semi Fowler's 27
 - side tilt 27
 - sitting 27
 - supine jack knife 28
 - 30 degree Trendelenburg 28
 - upper abdominal surgery 27 28
- Posterior fossa explorations 27
- Post operative care 105
- Postural ischemia 25 29
- Preganglionic sympathetic fibers
 - blocking of 22
 - procaine 43
- Pregnancy
 - hypertensive complications of 56
 - toxic states of 50

Tonsillectomy 94
 Total spinal analgesia 43 44
 Total spinal block
 basic features of 43 44
 in controlled hypotension
 advantages 45-46
 disadvantages 46

Tourniquets 11

Toxemia 6

Toxicity high degree of 5

Tracheotomy 85

Transfusion

 avoidance of 98

 reactions 98

Trendelenburg position 45 68

 reverse 27 45 68

 30 degree 28

Trichlorethylene 52

Tubular reabsorption 72

Tumors

 vascular 6

 brain 96

 removal of 31 94

U

Ulcer

 duodenal 50

 gastric 50

Ulcerative colitis sympathetic over
 activity 56

Urea 71

Urine formation 72

 suppression of 71

Urological surgery 96

V

Vascular bed

 capacity of following sympathet-
 ic blockade 26

 dilated 104

Vascular deficiency 5

Vascular disease 102

Vascular lesions removal of 31

Vascular ooze 3 7 9 11 46 58

 abnormal 5

 control of 14

Vascular sclerosis 51

Vascular tension 98

Vagal tone increase in 63

Vaginal plastic operation 96

Vasoconstriction 9 62 86

 agents 45 54

 arteriotomy 35

 fibers 40

 impulses 20 22

 interruption of 21 48

 peripheral 13 31

Vasopressor agents 54

 of Shorr 71

 reactionary hemorrhage 76

 use of 73 74

Vasodilatation 7 50 88 92

 and carbon dioxide absorption
 10

 during deep general anesthesia 8

 effect of muscle relaxants 9

 spinal analgesia 42

Vasomotor center

 depression 20

 in medulla oblongata 20

 reflex inhibition of 20

Veins small contraction of 20

Venodilatation 25 29

Venous congestion 5 9

Venous drainage 25 104

 gravitational 6 7 12

Venous ooze 10 12

Venous pressure 9 63

 increased 5

Venous return 17 63

 and continued circulation 29

 spinal analgesia 44-45

Ventilation 72

Ventricular failure acute 50

Veratrum alkaloids 22

Vesico-vaginal fistula 96

Vision blurred 77

Vital organs ischemia to 27 28

Y

Yawning 68

- oxygen saturation 65
- peripheral blood flow 66
 - low 66
- peripheral respiratory paralysis 42
- total 22 43 44
- vasodilatation 42
- Spinal anesthesia 12 14
 - high 81 86 87
- Spinal block 90
 - coronary flow 65
 - differential 42 68
 - total 14 41-46
 - in controlled hypotension 24
- Spinal fusion 96
- Splanchnic blood flow 66
- Splanchnic vasodilatation 62
- Subarachnoid block 42
- Succinylcholine and muscular relaxation 9
- Superior maxilla excision of spinal analgesia 41
- Surgery
 - bleeding during 3 5 10
 - and position of patient 6 7
 - anesthetic factors 7 10
 - control of 11 15
 - factors influencing 6
 - bloodless operative site 3-4
 - coagulation
 - effect of chloroform 7
 - effect of cyclopropane 7
 - effect of ether 7
 - effect of ethylene 7
 - effect of intravenous barbiturates 7
 - effect of nitrous oxide 7
 - emergency 98
 - head and neck 95
 - hemorrhage during 3
 - involving glandular tissue 6
 - involving muscles 6
 - Lempert fenestration 3
 - muscular relaxation 9
 - of the extremities 11
 - position of patient 12 14
 - and bleeding 6 7
 - uncomplicated and elective 89 92 95
 - vascular ooze 3
 - vasoconstrictor substances at operative site 11
- Sweating 92
- Sympathetic blockade 22 37 64 73
 - and capacity of vascular bed 26
 - severe hypotension 29
- Sympathetic chain block 40
- Sympathetic ganglia 48
 - blockade 22
- Sympathetic paralysis 22 70
- T
- Tachycardia 59 63 64
- Tachyphylaxis 59
- Tetraethylammonium bromide 22
- Thiopentham compound 55
- Thorax
 - pump action of negative pressure 25
 - wall lift of 9
- Thoracolumbar splanchnicectomy
 - position of patient 27
- Thoraco-lumbar sympathectomy 95
 - position of patient 27
- Thoracoplasty 95
- Thyroidectomy 6 95
 - spinal analgesia 41
 - subtotal for toxic goiter 84
- Thromboplastin deficiency 5
- Thrombosis 67
- Thrombotic phenomena 78 83
- Thrombus formation 29 88 89
- Tissue
 - anoxia 40
 - glandular surgery involving 6
 - oxygenation 104
 - level of pressure 40
 - vascularity of 6

